



COMPARATIVE STUDIES

CROSS NATIONAL SUMMARIES

NUMBER 9 – JUNE 1981

Contraceptive Practice

ENRIQUE CARRASCO

INTERNATIONAL STATISTICAL INSTITUTE
Permanent Office. Director: E. Lunenberg
428 Prinses Beatrixlaan
Voorburg, The Hague
Netherlands

WORLD FERTILITY SURVEY
Acting Project Director:
V. C. Chidambaram,
35–37 Grosvenor Gardens
London SW1W 0BS, U.K.

The World Fertility Survey (WFS) is an international research programme whose purpose is to assess the current state of human fertility throughout the world. This is being done principally through promoting and supporting nationally representative, internationally comparable, and scientifically designed and conducted sample surveys of fertility behaviour in as many countries as possible.

The WFS is being undertaken, with the collaboration of the United Nations, by the International Statistical Institute in co-operation with the International Union for the Scientific Study of Population. Financial support is provided principally by the United Nations Fund for Population Activities and the United States Agency for International Development. Substantial support is also provided by the UK Overseas Development Administration.

For information on Country Reports, WFS publications, and WFS depository libraries, write to the Publications Office, International Statistical Institute, 428 Prinses Beatrixlaan, PO Box 950, 2270 AZ Voorburg, Netherlands. For information on the WFS generally, write to the Information Office, World Fertility Survey, International Statistical Institute, 35-37 Grosvenor Gardens, London SW1W 0BS, UK.

L'Enquête Mondiale sur la Fécondité (EMF) est un programme international de recherche dont le but est d'évaluer l'état actuel de la fécondité humaine dans le monde. Afin d'atteindre cet objectif, des enquêtes par sondage sur la fécondité sont mises en oeuvre et financées dans le plus grand nombre de pays possible. Ces études, élaborées et réalisées de façon scientifique, fournissent des données représentatives au niveau national et comparables au niveau international.

L'EMF est entreprise, en collaboration avec les Nations Unies, par l'Institut International de Statistique, qui coopère avec l'Union internationale pour l'étude scientifique de la population. Le financement de ce programme est essentiellement assuré par le Fonds des Nations Unies pour les activités en matière de population et par l'Agence des Etats-Unis pour le développement international. Une contribution importante est aussi faite par le Département pour le développement des pays d'outre-mer du Royaume-Uni.

Pour toute information concernant les rapports d'enquêtes nationaux, les publications de l'EMF ou les bibliothèques dépositaires, écrire au Bureau des publications, Institut International de Statistique, 428 Prinses Beatrixlaan, BP 950, 2270 AZ Voorburg, Pays-Bas. Pour tous renseignements complémentaires sur l'EMF en général, écrire au Bureau d'information, Enquête Mondiale sur la Fécondité, Institut International de Statistique, 35-37 Grosvenor Gardens, Londres SW1W 0BS, Royaume-Uni.

La Encuesta Mundial de Fecundidad (EMF) es un programa internacional de investigación cuyo propósito es determinar el estado actual de la fecundidad humana en el mundo. Para lograr este objetivo, se están promoviendo y financiando encuestas de fecundidad por muestreo en el mayor número posible de países. Estas encuestas son diseñadas y realizadas científicamente, nacionalmente representativas y comparables a nivel internacional.

El proyecto está a cargo del Instituto Internacional de Estadística, contando con la colaboración de las Naciones Unidas y en cooperación con la Unión Internacional para el Estudio Científico de la Población. Es financiado principalmente por el Fondo de las Naciones Unidas para Actividades de Población y por la Agencia para el Desarrollo Internacional de los Estados Unidos. La Oficina Británica para el Desarrollo de Países Extranjeros proporciona también un gran apoyo financiero.

Puede obtenerse información sobre Informes de Países como otras publicaciones de la EMF y las bibliotecas depositarias, escribiendo a la Oficina de Publicaciones, Instituto Internacional de Estadística, Prinses Beatrixlaan 428, Casilla Postal 950, 2270 AZ Voorburg, Países Bajos. Si desea información de carácter general sobre la EMF, escriba a la Oficina de Información, Encuesta Mundial de Fecundidad, Instituto Internacional de Estadística, 35-37 Grosvenor Gardens, Londres SW1W 0BS, Reino Unido.

COMPARATIVE STUDIES

CROSS NATIONAL SUMMARIES

Contraceptive Practice

**ENRIQUE CARRASCO
WFS CENTRAL STAFF
International Statistical Institute
35-37 Grosvenor Gardens
London SW1W 0BS, UK**

Contents

PREFACE	5
ACKNOWLEDGEMENTS	6
INTRODUCTORY NOTE	7
1 VARIABLES	9
1.1 Contraceptive Use Variables	9
1.2 Demographic Variables	9
1.3 Control Variables	9
1.4 Content of the Tables	9
2 THE COLLECTION OF BASIC INFORMATION	10
2.1 The Individual Questionnaire	10
2.2 Specific Questions on Contraceptive Use	10
2.3 Variables Other than Use of Contraception	11
3 COMMENTARY ON THE DATA	12
3.1 Ever-Use of Contraception	12
3.2 Ever-Use of Individual Methods	13
3.3 Current Use of Contraception	16
3.4 Current Use of Individual Methods	18
REFERENCES	23
TABLES	
Group 1 Ever-Use by Type of Method	25
Group 2 Ever-Use by Individual Methods	29
Group 3 Current Use by Type of Method	43
Group 4 Current Use by Individual Methods	49
FIGURES	
1 Currently Married Women: Percentage of Ever-Use of Contraception, by Type of Method	13
2 Currently Married Women: Percentage of Ever-Use of Contraception, by Current Age	14
3 Currently Married Women: Percentage of Ever-Use of Contraception, by Number of Living Children	15
4 Exposed Women: Percentage of Current Use of Contraception, by Type of Method	17
5 Patterns of Current Use, by Age and by Number of Living Children	18
6 Exposed Women: Percentage of Current Use of Contraception, by Current Age	19
7 Exposed Women: Percentage of Current Use of Contraception, by Number of Living Children	20
8 Exposed Women: Percentage Using Modern Contraception	21

Preface

The first issues of the Cross National Summaries in the Comparative Studies series provide basic information, documentation and results of the World Fertility Survey for the nineteen countries which had their First Country Reports and Standard Recode Tapes available at the beginning of 1980.

Despite the efforts made by WFS to maintain comparability of question wording and content, field procedures and specifications of the tabulations and analysis included in the First Country Reports, it was inevitable that differences would arise as a result of the importance attached to meeting specific requirements of the countries themselves. A major attempt to enhance and facilitate comparability has been the production of Standard Recode Tapes for each country, with all the core information coded and stored in a consistent order, together with the dictionaries which provide detailed specifications for all variables.

Several of the Cross National Summaries will be concerned solely with providing detailed and systematized information on the comparability (or lack thereof) of the field procedures, survey characteristics, questionnaire content and wording and content of the First Country Reports. Such detailed appraisals constitute an essential reference base for anyone using WFS data for comparative analysis.

Other volumes of the Cross National Summaries will present comparable results from as many surveys as possible. These volumes will present the basic data from the surveys over a wide range of specific topics. In addition to the tabular material, there will be a brief accompanying text, which will draw attention primarily to any non-comparability of the data and to any obvious interpretational pitfalls to which the tables may be subject: for example many summary indices are subject to compositional differences, which are often reduced by standardisation. Finally, although these volumes are not intended to be analytic in their orientation, some brief highlighting of the major noteworthy differences and similarities is included.

We hope that these Cross National Summaries will be widely used especially by persons in the international community who are making cross national comparisons. We also hope that the sub-series will help users to avoid assuming too much comparability when this is not the case and to avoid interpretational mistakes which can easily arise when data are presented without qualification.

V.C. Chidambaram
Acting Project Director

Acknowledgements

Although authorship is attributed to the person(s) taking primary responsibility for the production of each of the Cross National Summaries, the work has been a co-operative effort involving many staff members of WFS. In particular, the production of the tables would often have been impossible without the substantial assistance of staff in the Data Processing Division and of Pat Platridis, Statistical Assistant.

The overall planning and co-ordination of the Cross National Summaries has been the responsibility of an editorial committee consisting of V. C. Chidambaram, John Cleland, John Hobcraft, Judith Rattenbury, German Rodriguez, Vijay Verma and Waller Wynne.

Introductory Note

This monograph presents comparative results of past and present use of contraception from studies in 19 countries, mainly by giving the percentages of women who ever used or currently use a contraceptive method. Our intention is not to present a comparative analysis of contraceptive use, which will be covered in future WFS publications, but rather to provide a description of similarities and differences in levels and patterns of use. For this reason contraceptive use, demographic and control variables have been included, but background variables excluded. It should be noted that a monograph dealing with urban-rural differences in use of contraception has also been published in this series (Lightbourne 1980). Many analysts will be interested in knowledge of contraception, and might like to know that this monograph and Vaessen's "Knowledge of Contraceptive Methods" (1980) were prepared using a similar presentation, enabling the two studies to be compared.

1 Variables

The types of variables employed here, namely use of contraception, demographic and control, are discussed below.

1.1 CONTRACEPTIVE USE VARIABLES

The principle aspect of measuring contraceptive use is timing. Accordingly, two variables are used: ever-use, which implies use at any time, for however long; and current use, with no distinction made between prolonged use and recent adoption. These variables are discussed further in chapter 2.

Another important aspect is the efficiency of a particular contraceptive method. Where figures are given for individual methods, the efficiency is immediately apparent. Elsewhere, a twofold classification is followed: efficient methods, ie sterilization (both male and female), pill, injection, IUD (any kind), condom and "other female scientific", such as diaphragm, jelly, foam, sponge, vaginal tablets and spermicides in general; and inefficient methods, ie rhythm, douche, withdrawal, abstinence and the wider category of folk methods. The distinction between efficient and inefficient methods is merely conventional and aims at distinguishing the more efficient from the less efficient. No attempt was made during the interview to assess whether the respondent used the method efficiently, and it should be noted that an efficient method may be used inefficiently and vice versa, though this is less common.

Sterilization, of course, differs from most contraceptive methods in being non-reversible, an important distinction. Nevertheless it shares with other methods the practical effect of preventing conception. Moreover, many women mention sterilization as a contraceptive method in the same way that they mention abortion when asked about contraception. The questionnaires have a series of questions which distinguish women sterilized for contraceptive purposes from those sterile for other reasons, and the term sterilization should be understood here as meaning "sterilized for contraceptive purposes".

1.2 DEMOGRAPHIC VARIABLES

The current age of respondents and the number of living children are employed as demographic variables in the study of contraceptive practice. Using age as a variable enables us to relate contraceptive practice to the physiological aspects of human reproduction and to social aspects, such as marriage, separation and divorce. Furthermore family planning programmes usually keep records of acceptors by age, providing a basis for comparison. In the present study, the current age of respondents is classified in five-year groups, from 15-19 to 45-49.

Although age is a commonly used demographic variable, a more important factor in the study of contraception is the number of living children, ie the number of children born to the women interviewed who are currently alive, whether or not they reside with her. Analysis of the relationships between contraception and background variables (eg socio-economic and cultural variables) should include comparisons of women with the same number of living children.

1.3 CONTROL VARIABLES

These variables refer to the population for whom the ever-use or current use of contraception is relevant. In the preparation of the country reports, this population has been referred to as the base of the tables. Four bases have been used in this monograph: ever-married; currently married; fecund, ie currently married women who believe themselves physiologically capable of having children, or pregnant, or sterilized for contraceptive purposes; and exposed, ie all fecund women except those who are pregnant.

Only minor differences in contraceptive practice are to be expected between ever-married and currently married women, and between fecund and exposed women. Since the results confirm this expectation, the tables on individual methods are limited to currently married women for ever-use (group 2) and currently married and exposed women for current use (group 4).

1.4 CONTENT OF THE TABLES

Each table shows the proportion of women who ever used or are using contraceptives, cross-classified by one of the demographic variables and confined to the specific base. In the case of currently married women using a contraceptive method, the percentages may be compared with the estimates of coverage provided by family planning programmes. These coverage estimates do not always refer to ever-married or currently married women, the most common exception being percentages referring to all women in the fertile age range. In this case, an adequate comparison can be made by using the proportion, either ever or currently married, of all women in the specific group to adjust the figures presented here (the proportions can be obtained from the First Country Report of each national fertility survey).

2 The Collection of Basic Information

2.1 THE INDIVIDUAL QUESTIONNAIRE

The national studies developed under the WFS programme are based on a household and an individual survey. The information collected in the household schedule was used, among other things, to identify women eligible for the main interview, which used the individual questionnaire. The eligible women are usually defined as those ever married under 50, or, in most Latin American and Caribbean countries, all women aged 15-49. For further details of the characteristics of the surveys, see Singh (1980a). Information on knowledge and use of contraception can be obtained from the results of the individual questionnaire, from section 3, Contraceptive Knowledge and Use, section 5, Fertility Regulation, and the Family Planning Module, which, added to section 3, has been used in some countries. Of the 19 countries considered in this monograph, only three (Colombia, Dominican Republic and Mexico) asked single women these questions. The present comparative results are based on the replies of women aged 15-49 who had ever been in a union, with the exception of Costa Rica and Panama, where women aged 15-19 were not interviewed. Further information on the comparability of questionnaires can be found in Singh (1980b).

2.2 SPECIFIC QUESTIONS ON CONTRACEPTIVE USE

The following comments are based on the English version of the individual core questionnaire recommended by the WFS. Knowledge and ever-use of contraception were ascertained in section 3. Question 301 on knowledge ("As you may know, there are various ways that a couple can delay the next pregnancy or avoid pregnancy. Do you know of, or have you heard of, any of these ways or methods?") was followed by the open-ended question 302 ("Which methods do you know of?"). All countries except Fiji recorded methods spontaneously mentioned in reply to this question, and in the case of Fiji, the exception does not seem to have affected the comparability of the information. All except Pakistan then probed for knowledge of each method not spontaneously mentioned, and for ever-use of all known methods. Hence the results for Pakistan are not comparable because the extent of use of contraceptive methods is under-reported. The national decision not to probe and to ask about ever-use only of those methods which were spontaneously mentioned was based on the findings of previous KAP surveys, which suggest that knowledge of contraceptive methods is overestimated when probes are used. However, WFS figures suggest that without probes under-reporting of knowledge would be similar or greater (see Vaessen, forthcoming).

Table I summarizes the comparability of information on knowledge and use of contraception. The quality of the information on use of contraception is affected by the procedure used to ask about knowledge. Both over and under-reporting are likely to occur, a topic discussed elsewhere in this series (Vaessen 1980).

Overestimating knowledge may be due, in part, to the

interviewer forcing a reply or the tendency of the respondent to answer yes, as a form of polite social response (known as the "social acquaintance response"). Unfortunately it is impossible to assess the extent of overestimation, but it may be considered less relevant to the use of contraception because inaccurate reporting of knowledge of a contraceptive method does not imply that the method is also used.

Table I Factors Affecting Comparability of Contraceptive Use Information

Country	No Probe on Knowledge	Use Question Not Asked	Age Group Not Included
<i>Asia and Pacific^a</i>			
Bangladesh	Injection	Injection	—
Fiji	Douche	Douche ^b	—
Indonesia	—	—	—
Jordan	Injection	Injection	—
Korea, Rep. of	—	—	—
Malaysia	Douche	Douche	—
Nepal	Injection, Rhythm, Other Female Scientific, Douche, Withdrawal	Same	—
Pakistan	All Methods	Douche ^c	—
Philippines	Injection	Injection	—
Sri Lanka	—	—	—
Thailand	—	—	—
<i>Caribbean and Latin America</i>			
Colombia	—	—	—
Costa Rica	—	—	15-19
Dominican Rep.	Abstinence	Abstinence	—
Guyana	—	—	—
Jamaica	—	—	—
Mexico	Abstinence	Abstinence	—
Panama	—	—	15-19
Peru	—	—	—

^a Including West Asia.

^b Included in "other female scientific".

^c Not coded as specific method.

Underestimating knowledge is more serious, and is probably mainly due to the inability of the interviewer to probe and the failure to give the respondent enough time to answer the open-ended question on knowledge.

In later sections of the individual questionnaire (section 5, Fertility Regulation, or the Fertility Regulation Module,

whichever was used), the interviewer was sometimes able to uncover ever-use or current use of a contraceptive method, which had not been reported in section 3, either spontaneously or after probing, thus discovering knowledge of a method which had not been mentioned earlier. However, overestimation of knowledge cannot be brought to light in the same way.

Whether the Family Planning Module was used is not relevant to a study of the use of contraception because women who knew about the supply of contraceptives (pill, IUD, condom, sterilization) were only asked about the outlets, accessibility and cost of supplies in this module. Current use of a method was ascertained in section 5 of the individual core questionnaire or in the Fertility Regulation Module, which may replace section 5, adding further questions and not omitting any. In any case, the questions on current use are not affected by the variations.

Through filter questions at the beginning of the section or module, it was possible to classify each woman as follows:

- (i) married and living with husband;
- (ii) not currently pregnant (or does not know);
- (iii) has used a contraceptive method.

In case of affirmative answers for each one of the filters, question 504 was asked ("Are you or your husband currently using a method to keep you from getting pregnant?"), and then question 505 ("What method are you using?"). In the case of sterilization, question 526 of section 5 or question 572 of the module ask if one purpose of that operation was "to prevent (her) having any children" thus permitting the category "woman (or husband) sterilized for contraceptive reasons" to be constructed.

2.3 VARIABLES OTHER THAN USE OF CONTRACEPTION

The current age of respondents was investigated in section 1, by means of question 107 ("In what month and year were you born?") or question 108 ("How old are you?"), the instruction for the latter being to record the best estimate. The intention in this paragraph is not to discuss the quality of this variable (this matter has been investigated extensively by country researchers and WFS staff and will be discussed in the forthcoming Data Evaluation Reports) but simply to point out that a five-year age group is appropriate for study of the general level and pattern of use of contraception, as well as for comparing figures among countries. An additional way of taking age into account is to standardize by age distribution, and the Fiji age distribution has been used for this purpose.

Although in some countries (Bangladesh, Republic of Korea, Malaysia, Pakistan, Sri Lanka, Thailand) women under 15 were included in the survey, and women of 50 interviewed in some of these and in other countries (Indonesia, Republic of Korea, Pakistan), the present report covers only women aged 15-49. In Costa Rica and Panama, women 15-19 were not eligible for the individual interview at all.

The number of living children (ie the number of children born to the women, currently alive, living or not with her) is provided by the maternity history (section 2), where a special effort was made to obtain the total number of children born and those still living. The distribution for Fiji was applied to obtain standardized use percentages.

Information on marital status was obtained from the marriage history (section 4). Probing, in order to determine the current marital status in the most accurate way, was employed. In Latin American and Caribbean countries, the ever-married status must be understood as "ever in union", since consensual unions were included. The category of "visiting union" was included for Guyana and Jamaica as well.

Question 221 ("Are you pregnant now?") identifies the current pregnancy status; the "don't know" answers were added to the category of "non-pregnant". The response to the question "As far as you know, is it physically possible for you and your husband to have a child, supposing you wanted one?", used jointly with currently pregnant and currently married information enables us to identify the fecund and exposed groups of women. The fecund group consists of women currently married who believe themselves physiologically capable of having children, and those who are pregnant or have been sterilized for contraceptive reasons. The exposed group is similar to the fecund group, but currently pregnant women are excluded.

Both for the currently pregnant and currently fecund status, it is generally felt that under-reporting exists (see Ferry 1980 and forthcoming; and Hanenberg 1980). The underestimate of the proportion of women currently pregnant is evident from a comparison with the general fertility rate or from the low percentages reporting pregnancies of three months or less. Of the women classified as fecund, an undetermined proportion could actually be infecund because of unrealized sterility, especially among older women. For the purposes of this study, it is assumed that the women who inaccurately report that they are not pregnant are distributed between current users and non-users in the same way as those who are genuinely not pregnant. The same assumption applies to the group inaccurately identified as fecund.

3 Commentary on the Data

All the countries discussed in this series of Cross National Summaries have already published their First Country Reports or will soon do so. Any differences in the figures in these reports and the present monograph may be attributed to further editing of the data-file at the WFS headquarters, but the discrepancies are not large and can be ignored.

Data were arranged into four main groups of tables: ever-use of contraception (any method), ever-use of individual methods, current use of contraception (any method) and current use of individual methods. The variables involved in each group of tables are presented as follows:

Use Variables	Demographic Variables	Control Variables	No of Tables
1 Ever-use by type of method	Current age, number of living children	Ever married, currently married	4
2 Ever-use by individual methods	Current age, number of living children	Currently married	24
3 Current use by type of method	Current age, number of living children	Ever married, currently married, fecund, exposed	8
4 Current use of individual methods	Current age, number of living children	Currently married, exposed	48

The following conventions for the presentation of the tables have been observed:

- Countries are arranged in alphabetical order within the respective region, using the term "region" for comparative purposes only.
- No decimal places are shown for percentages in tables of groups 1 and 3.

3.1 EVER-USE OF CONTRACEPTION

Tables in group 1 show the percentages of women who ever used contraception, both for ever-married and for currently married women. As might be expected, the percentages for the latter group are lower, but the patterns remain similar. Age-standardized values are also presented, but these values do not differ significantly from the observed ones. The differences between observed values and values standardized by age or by number of living children are small. Note that the correction introduced by the standardization may have one of two completely opposite effects, depending on whether the factor of age or number of living children is taken into account. This is not always the case with standardization of current use. Therefore, the following comments refer exclusively to observed values for currently married women (tables 1.2.1 and 1.2.2).

The general level of ever-use is shown in figure 1, where the countries are arranged, within each region, according to the total fertility rate (from high to low) estimated for

the three years preceding the survey. The level of use varies extensively, from 5 and 15 per cent in Nepal and Bangladesh respectively to 86 per cent in Costa Rica.

Generally speaking, the Latin American and Caribbean countries reveal higher percentages of ever-use than the Asian and Pacific countries. Although contraceptives can be obtained with or without the assistance of family planning programmes, these programmes began earlier in Latin America and the Caribbean and are already highly developed in several countries. With the exception of Peru, the ever-use of efficient methods is always surprisingly higher than the use of inefficient methods. Where the level of ever-use remains low, one might also expect a relatively low prevalence of efficient methods (sterilization, pill, IUD, injection, condom, other female scientific). However, figure 1 suggests that in the Asian and Pacific countries, contraceptive practice is skipping the stage where inefficient methods predominate. Ten or fifteen years ago, when the first KAP studies were carried out in Latin America and the Caribbean, a considerable amount of ever-use consisted of inefficient methods.

As explained earlier, the countries in figure 1 are arranged according to the total fertility rate estimated for the last three years, and it should be noted that a high level of ever-use is not always associated with low fertility. The total fertility rates are as follows (see Hanenberg 1980).


Mexico	5.7	Colombia	4.4
Dominican Republic	5.3	Guyana	4.4
Peru	5.2	Panama	4.0
Jamaica	4.5	Costa Rica	3.5
Jordan	7.0	Malaysia	4.3
Bangladesh	6.1	Indonesia	4.2
Nepal	6.0	Thailand	4.2
Pakistan	5.8	Korea, Rep. of	4.0
Philippines	4.8	Fiji	3.9
		Sri Lanka	3.4

Figure 1 shows clearly that in populations of similar fertility levels, completely different contraceptive behaviour can be reported, especially in the Asian and Pacific countries. The contrast, for instance, between Jordan, on the one hand, and Bangladesh, Nepal and Pakistan, on the other, is notable.

Figures 2 and 3 show proportions of currently married women who have ever used a contraceptive method, by age or number of living children. These percentages do not indicate when the women used contraception for the first time, nor how frequent use of contraception was. Because of the cumulative nature of the category of ever-use, women who once used a contraceptive method many years ago are included in the category, along with regular users at the time of the interview.

The following comments refer only to the patterns shown by numbers of living children. Figure 3 shows several different patterns. One of them, revealing high and well-

Figure 1 Currently Married Women: Percentage of Ever-Use of Contraception, by Type of Method



spread contraceptive practice, is that of Costa Rica and Panama. For these two countries one must bear in mind that the figures are calculated for women aged 20-49 only, and lower percentages (mainly for zero, one and two living children) could be assumed if women aged 15-19 were included.

It is striking that several developing countries are already exhibiting a pattern similar to that of developed countries. Costa Rican and Panamanian women with only one living child show the same percentage using contraception (90 and 80 per cent respectively) as women with two, three or four living children. Colombia, Jamaica and Fiji also seem to be approaching this pattern of earlier use.

A wholly different pattern is that of Bangladesh, Nepal and Pakistan, where ever-use is rare. The figures suggest that women have begun to use contraceptive methods only after having a considerable number of children, showing a pattern of later use. This pattern also prevails in Republic of Korea, though contraceptive use is considerably higher.

3.2 EVER-USE OF INDIVIDUAL METHODS

The tables in group 2 show the ever-use of individual methods for currently married women. The same woman

Figure 2 Currently Married Women: Percentage of Ever-Use of Contraception, by Current Age





Figure 3 Currently Married Women: Percentage of Ever-Use of Contraception, by Number of Living Children



may report using more than one method. Table II includes only the methods most frequently mentioned.

PILL . This is the most frequently reported method in every country with the exception of the Philippines, Sri Lanka and Peru. Despite its popularity, the extent of its use varies considerably.

IUD This method is substantially less frequently reported than the pill. Republic of Korea (26 per cent), Fiji (17 per cent), Colombia (16 per cent) and Costa Rica (13 per cent) have the highest percentages.

FEMALE STERILIZATION FOR CONTRACEPTIVE PURPOSES These percentages are among the lowest values, especially in the Asian and Pacific countries. Panama (21 per cent) and Fiji (16 per cent) have the highest percentages. The proportion of currently married women who are sterilized (table 2.9.2) rises sharply for women with three or more living children in Panama (26 per cent) and for those with four or more living children in Fiji (23 per cent).

MALE STERILIZATION This is rarely reported, and the only country worth noting is Republic of Korea with 3.3 per cent (table 2.10.1).

CONDOM The condom has been widely used in a high proportion of the countries. Costa Rica (37 per cent), Jamaica (26 per cent), Republic of Korea (22 per cent) and Philippines (21 per cent) show the highest values.

RHYTHM AND WITHDRAWAL These are reported as frequently as the condom, with Philippines, Peru, Costa Rica, Colombia, Panama and Fiji showing the greatest ever-use. In Philippines and Peru these two methods are the most frequently ever used.

OTHER FEMALE SCIENTIFIC This group consists of mechanical or chemical barriers. Reported use is not significant in the Asian and Pacific countries, but in the Latin American and Caribbean countries ever-use is substantial.

3.3 CURRENT USE OF CONTRACEPTION

This information is given in the tables of group 3, for ever-married, currently married, fecund and exposed women. The percentages are very similar for ever and currently married women, on the one hand, and for fecund and exposed women, on the other. Once again the observed and standardized values do not differ (indeed the discrepancy is even less than in the case of ever-use). The commentary is therefore confined to observed values for exposed women, defined as women currently married and not pregnant who believe themselves capable of having children, plus those sterilized for contraceptive reasons.

General Level of Current Use

The general level of current use is illustrated in figure 4, where the countries have been arranged within each region according to the total fertility rate (from high to low) estimated for the three years preceding the survey.

Broadly speaking, the observations made for ever-use are valid here. There is, most notably, a wide variation in the percentages, ranging from as low as 10 per cent or less in Nepal, Pakistan and Bangladesh to 65 per cent in Panama and 78 per cent in Costa Rica. Different per-

Table II Percentage of Currently Married Women Who Ever Used an Individual Contraceptive Method

Country	Contraceptive Method						
	Pill	IUD	Female Sterilization ^a	Condom	Rhythm	Withdrawal	Other Female Scientific ^b
<i>Asia and Pacific^c</i>							
Bangladesh	5.9	1.2	0.3	2.3	5.5	3.0	0.6
Fiji	31.6	16.5	15.8	16.7	15.8	15.5	3.5
Indonesia	26.2	8.0	0.3	5.4	3.0	2.1	0.3
Jordan	36.5	4.5	1.8	8.9	13.2	18.9	1.7
Korea, Rep. of	35.0	25.7	1.7	22.5	16.8	12.3	0.6
Malaysia	34.1	1.6	3.4	9.1	11.0	8.3	1.8
Nepal	2.0	0.2	0.1	0.7	N Ask	N Ask	N Ask
Pakistan	3.7	3.1	0.9	2.6	0.2	0.3	0.9
Philippines	25.2	7.2	4.7	20.7	23.5	31.7	3.0
Sri Lanka	8.7	8.1	9.2	5.1	22.5	6.5	0.3
Thailand	27.3	9.9	6.1	3.6	7.3	5.4	1.0
<i>Caribbean and Latin America</i>							
Colombia	34.5	16.3	4.0	8.8	19.7	18.1	12.5
Costa Rica ^d	53.0	12.9	12.3	37.0	21.7	26.8	12.0
Dominican Rep.	21.3	6.9	11.9	10.3	7.3	19.0	7.2
Guyana	21.8	8.8	8.5	16.5	11.1	13.9	8.8
Jamaica	31.1	8.1	8.1	25.6	5.2	20.6	11.2
Mexico	26.7	8.6	2.7	6.6	13.3	18.3	8.0
Panama ^d	44.0	9.1	21.2	13.3	16.4	18.5	9.9
Peru	3.9	2.7	2.8	7.0	27.4	16.8	5.0


^a For contraceptive purposes only.

^b Diaphragm, jelly, foam, sponge, vaginal tablets, spermicides in general.

^c Including West Asia.

^d Women aged 20-49 only.

Figure 4 Exposed Women: Percentage of Current Use of Contraception, by Type of Method



centages of current use in countries with similar levels of fertility should also be noted, and the use of efficient methods is greater than the use of inefficient ones, with the exception of Peru and Philippines.


If the level of current use is compared with that of ever-use, it is evident that the percentages are closely correlated. The level of current fertility in two countries (Guyana and Sri Lanka) is seemingly inconsistent with their contraceptive practice, expressed either as ever-use or current use, suggesting that a factor other than contraception is strongly affecting fertility.

Two other countries (Peru and Philippines) deserve special mention in a comparison of past and present use of efficient methods. In both countries women show high current levels of use of inefficient methods, with Peru maintaining the same pattern of behaviour as it has had in the past and Philippines showing a change of direction.

Patterns of Current Use by Current Age and Number of Living Children

These patterns are shown in figures 6 and 7, which are based on tables 3.4.1 and 3.4.2. In figure 5, three different patterns are shown (they are the smoothed values of Nepal,

Figure 5 Patterns of Current Use by Age and by Number of Living Children
(Three Typical Observed Situations)



Dominican Republic and Costa Rica) with the aim of enabling comparison between countries in figures 6 and 7. Curves (or shapes) of type C represent both low current use and very recently developed contraceptive practice or family planning programmes. Contraception is used mainly by older women with a large number of living children. At the other extreme, type A curves represent high widespread incidence of contraception. Contraceptive methods are used at earlier stages of the reproductive period by women of younger ages with a small number of living children. The grounds for contraceptive practice may be expressed as "spacing" and "limiting". An intermediate situation is found in the curves of type B, where the maximum values of contraceptive use are as likely to occur at the older or middle ages and at the highest number of children or at more moderate values.

Countries like Costa Rica and Panama may be classified as type A, while Nepal, Pakistan and Bangladesh belong to type C. In the countries classified as type B, some differential patterns can be distinguished according to whether the maximal values appear later (Fiji, Republic of Korea, Jordan) or earlier (Colombia, for instance). One notable pattern is that of Republic of Korea, where current use is comparatively low for women of younger ages or for those with zero or one living child and then rises sharply. In terms of number of children, it seems that Korean women are not concerned with contraception while they have zero or only one child, but after the second, use of contraception becomes significantly higher.

The incidence of inefficient methods is clearly lower, with the exception of Peru and Philippines. The important fact to note here is that in both these countries the higher incidence of inefficient methods prevails at every age and in every number of living children group.

3.4 CURRENT USE OF INDIVIDUAL METHODS

These percentages by age and number of living children are presented in the tables of group 4, both for currently married and for exposed women. In table III, the overall percentages (all ages, or all living children) of current use in exposed women for each method are shown.

Modern Methods (Pill, IUD, Sterilization)

Because of low levels of use, injection is included with the pill and male sterilization is included with female sterilization.

The pill is one of the most frequently reported methods currently used, though not the most frequent in every country. The percentage of exposed women using the pill is very high in several countries (30 per cent in Costa Rica and 21 per cent in Thailand, Indonesia and Jamaica). Sterilization is more widely used than the pill in Panama (26 per cent), Fiji (20 per cent) and Dominican Republic (16 per cent). Of the modern methods, the IUD is never the most common, but its level of use is notable in Republic of Korea (11 per cent), Colombia (10 per cent), Indonesia and Mexico (8 per cent each).

The level of current use of modern methods is high in many countries, as shown in figure 8, where the overall level of current use for every country is given in brackets.

The pattern of use of modern methods by age and number of living children is very much what one might expect. For the pill the highest values are observed for younger women or women with a small number of living children, followed by monotonically declining percentages. The use of the IUD becomes significant when the number of

Figure 6 Exposed Women: Percentage of Current Use of Contraception, by Current Age




Figure 7 Exposed Women: Percentage of Current Use of Contraception, by Number of Living Children





Figure 8 Exposed Women: Percentage Using Modern Contraception



Note: General level of current contraceptive use is shown in parentheses.

Table III Percentage of Exposed Women Currently Using Contraception, by Individual Method Used

Country	Contraceptive Method							
	Pill ^a	IUD	Sterilization ^b	Any Modern	Condom	Rhythm	Withdrawal	Others
<i>Asia and Pacific^c</i>								
Bangladesh	3.7	0.5	1.0	5.2	0.9	1.4	0.6	1.8
Fiji	10.9	6.0	20.2	37.1	7.7	3.0	3.5	0.9
Indonesia	21.1	7.9	0.4	29.4	2.6	1.2	0.5	3.3
Jordan	17.6	3.0	2.8	23.4	2.1	3.1	4.9	3.8
Korea, Rep. of	11.3	10.5	6.6	28.4	6.7	6.0	3.4	1.2
Malaysia	21.0	0.9	4.9	26.8	3.5	4.4	2.5	5.2
Nepal	0.5	0.1	2.1	2.7	0.3	N Ask	N Ask	0.1 ^d
Pakistan ^e	1.3	0.9	1.4	3.6	1.4	0.2	0.1	2.0
Philippines	6.2	3.1	7.1	16.4	4.7	11.3	12.5	2.9
Sri Lanka	2.4	6.2	12.9	12.9	3.0	10.4	1.9	4.6
Thailand	21.3	8.1	11.5	40.9	0.6	1.2	1.2	1.4
<i>Caribbean and Latin America</i>								
Colombia	16.7	10.4	5.1	32.2	2.1	6.2	5.8	5.5
Costa Rica ^f	29.6	6.3	16.1	52.0	10.7	6.2	5.6	3.6
Dominican Rep.	10.5	3.7	15.7	29.9	2.0	1.6	4.9	3.2
Guyana	10.6	6.5	11.5	28.6	0.0	3.4	1.2	4.4
Jamaica	21.0	2.3	11.4	34.7	7.6	0.4	1.7	2.3
Mexico	17.2	7.8	4.0	29.0	1.0	4.2	5.0	2.4
Panama ^f	21.4	4.5	26.0	51.9	1.5	3.1	3.6	4.2
Peru	6.9	1.8	3.6	12.3	1.4	14.3	4.3	9.1

^a Includes injection.

^b Male and female sterilization for contraceptive purposes only.

^c Including West Asia.

^d Presumably underestimated.

^e Not strictly comparable.

^f In respect of women aged 20-49 only.

living children is two or more. Sterilization shows different patterns in different countries: in Panama and Dominican Republic, the incidence is considerable among exposed women with three living children (31 and 21 per cent), although figures for women with two living children are also substantial (12 and 10 per cent). In Fiji, Costa Rica, Sri Lanka and Thailand, female sterilization is widely used by women with four or more living children, and in Guyana and Jamaica by women with five or more living children.

Other Methods

Generally speaking, the incidence of methods other than the modern ones defined above is not of great significance, but a few should be mentioned here. In Philippines, rhythm and withdrawal and in Peru, rhythm alone were the most widely used methods at the time of the interview. In Costa Rica, the total percentage using non-modern methods is 26.1, indicating that both modern and traditional methods are widely used. A large proportion of women using traditional methods were among the older cohorts.

References

Ferry, Benoît (1980). Breastfeeding. *WFS Comparative Studies* no 13.

Ferry, Benoît (forthcoming). Some Factors other than Contraception Affecting Fertility. *WFS Scientific Reports*.

Hanenberg, Robert (1980). Current Fertility. *WFS Comparative Studies* no 11.

Lightbourne, Robert (1980). Urban-Rural Differentials in Contraceptive Use. *WFS Comparative Studies* no 10.

Singh, Susheela (1980a). Characteristics of the Surveys. *WFS Comparative Studies* no 1.

Singh, Susheela (1980b). Comparability of Questionnaires. *WFS Comparative Studies* no 2.

Vaessen, Martin (1980). Knowledge of Contraceptive Methods. *WFS Comparative Studies* no 8.

Vaessen, Martin (forthcoming). Knowledge of Contraception: an Assessment of WFS Data Collection Procedures. *Population Studies*.

Tables

Group 1 Ever-Use by Type of Method	25
Group 2 Ever-Use by Individual Methods	29
Group 3 Current Use by Type of Method	43
Group 4 Current Use by Individual Methods	49

NOTES

In the following tables Asia and Pacific include West Asia.

- a. Methods are grouped into two types: efficient (male and female sterilization, pill, IUD, injection, condom and other female scientific) and inefficient. Figures are given for three groups: None (N), Inefficient only (I) and some Efficient (E).
- b. According to Fiji age distribution.
- c. Only asked of women who spontaneously mentioned this method in answer to the question on knowledge of contraception.
- d. According to Fiji living children distribution.
- e. Figures for women aged 20-49 only.
- f. Including douche.
- g. Women who believe themselves physiologically capable of having children, or pregnant, or sterilized for contraceptive purposes.
- h. Women currently married, not pregnant who: a) believe themselves physiologically capable of having children, or b) have been sterilized for contraceptive purposes.

**The following abbreviations
have been used in the tables**

NA Not available

N Ask Not asked

Contraceptive methods:

N None

I Inefficient only

E Efficient

GROUP 1 EVER-USE BY TYPE OF METHOD

LIST OF TABLES

- | | | |
|-------|--|----|
| 1.1.1 | Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Ever Used, by Current Age | 27 |
| 1.1.2 | Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Ever Used, by Number of Living Children | 27 |
| 1.2.1 | Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Ever Used, by Current Age | 28 |
| 1.2.2 | Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Ever Used, by Number of Living Children | 28 |

Table 1.1.1 Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Ever Used, by Current Age

Country	Current Age and Type of Method ^a Ever Used														Total												
	15-19			20-24			25-29			30-34			35-39			40-44			45-49			Observed		Standardized ^b			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																											
Bangladesh	93	4	3	85	7	8	85	6	9	80	9	11	81	8	11	85	7	8	91	6	3	86	6	8	84	6	10
Fiji	59	10	31	39	10	51	27	9	64	22	12	66	28	11	61	32	14	54	48	13	39	32	11	57	32	11	57
Indonesia	84	2	14	65	3	32	54	4	42	55	5	40	60	5	35	71	5	24	82	6	12	66	4	30	63	4	33
Jordan	83	5	12	61	9	30	48	9	43	44	8	48	47	8	45	48	5	47	58	6	36	54	7	39	52	7	40
Korea, Rep. of	83	4	13	70	4	26	46	7	47	31	6	63	26	4	70	39	4	57	65	3	32	43	5	52	47	5	48
Malaysia	76	8	16	56	10	34	43	10	47	42	10	48	49	9	42	54	11	35	73	11	16	52	10	38	51	11	38
Nepal	99	0	1	98	1	1	95	1	4	93	1	6	92	1	7	95	1	4	97	1	2	96	1	3	95	1	4
Pakistan ^c	99	0	1	96	0	4	90	1	9	86	2	12	83	2	15	86	3	11	89	2	9	90	2	8	89	2	9
Philippines	74	13	13	49	16	35	35	19	46	33	17	50	34	19	47	45	19	36	63	17	20	42	18	40	42	18	40
Sri Lanka	81	9	10	68	12	20	57	17	26	45	20	35	48	19	33	57	20	23	66	22	12	57	18	25	57	18	25
Thailand	74	6	20	58	6	36	44	6	50	44	7	49	46	4	50	60	7	33	75	9	16	54	7	39	53	6	41
<i>Caribbean and Latin America</i>																											
Colombia	56	10	34	38	14	48	34	10	56	28	11	61	38	10	52	48	15	37	62	16	22	41	12	47	39	12	49
Costa Rica	NA	NA	NA	15	6	79	11	7	82	15	7	78	16	9	75	20	12	68	41	13	46	18	9	73	18	8	74
Dominican Rep.	70	12	18	53	14	33	39	10	51	44	11	45	47	9	44	57	11	32	70	9	21	52	11	37	50	11	39
Guyana	61	11	28	52	10	41	39	10	51	33	7	60	43	8	49	50	8	41	56	13	31	46	9	44	45	9	46
Jamaica	41	11	48	30	11	59	23	8	69	24	9	67	33	8	59	45	7	48	54	14	32	34	10	56	32	9	58
Mexico	74	7	19	56	9	35	44	10	46	46	10	44	51	11	38	61	10	29	72	10	18	55	10	35	54	10	37
Panama	NA	NA	NA	28	11	61	21	10	69	25	9	66	26	11	63	31	10	59	38	14	48	27	10	63	27	10	63
Peru	68	22	10	51	29	20	43	29	28	41	26	33	51	23	26	55	26	19	67	19	14	51	26	23	50	26	24

Table 1.1.2 Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Ever Used, by Number of Living Children

Country	Number of Living Children and Type of Method ^a Ever Used														Total									
	None			1			2			3			4			5+			Observed		Standardized ^d			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																								
Bangladesh	96	2	2	90	5	5	87	6	7	84	8	8	84	8	8	79	8	13	86	6	8	85	6	9
Fiji	75	9	16	44	14	42	30	12	58	26	11	63	22	9	69	19	11	70	32	11	57	32	11	57
Indonesia	95	1	4	74	4	22	59	5	36	55	5	40	56	5	39	54	5	41	66	4	30	62	5	33
Jordan	87	6	7	68	7	25	56	7	37	51	9	40	51	7	42	46	7	47	54	7	39	57	7	36
Korea, Rep. of	81	4	15	73	5	22	41	7	52	31	5	64	29	4	67	37	3	60	43	5	52	48	4	47
Malaysia	85	7	8	61	9	30	49	12	39	47	11	42	47	11	42	47	9	44	52	10	38	54	11	35
Nepal	99	1	0	98	1	1	97	1	2	95	1	4	92	1	7	90	1	9	96	1	3	94	1	5
Pakistan ^c	99	0	1	97	1	2	94	2	4	91	1	8	87	2	11	80	3	17	90	2	8	89	2	9
Philippines	88	6	6	52	20	28	41	20	39	32	20	48	34	18	48	41	18	41	42	18	40	46	17	37
Sri Lanka	91	7	2	69	18	13	53	25	22	49	22	29	44	20	36	50	17	33	57	18	25	57	18	25
Thailand	84	5	11	59	7	34	49	7	44	45	6	49	45	7	48	55	6	39	54	7	39	56	6	38
<i>Caribbean and Latin America</i>																								
Colombia	66	10	24	45	11	44	33	13	54	35	11	54	33	14	53	42	12	46	41	12	47	42	12	46
Costa Rica ^e	54	5	41	16	7	77	11	8	81	12	10	78	14	9	77	21	10	69	18	9	73	20	8	71
Dominican Rep.	75	11	14	62	10	28	42	13	45	42	12	46	41	10	49	50	11	39	52	11	37	51	11	38
Guyana	63	10	27	56	11	33	43	9	48	42	10	48	40	9	51	40	8	52	46	9	44	46	9	45
Jamaica	51	11	38	42	11	47	28	10	62	30	8	62	27	12	61	29	8	63	34	10	56	33	10	57
Mexico	81	6	13	58	9	33	44	11	45	48	9	43	50	11	39	56	10	34	55	10	35	56	10	35
Panama ^e	57	6	37	32	8	60	21	10	69	21	9	70	20	10	70	29	12	59	27	10	63	29	10	60
Peru	78	13	9	55	28	17	43	27	30	43	27	30	49	26	25	55	25	20	51	26	23	54	25	22

Table 1.2.1 Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Ever Used, by Current Age

Country	Current Age and Type of Method ^a Ever Used												Total														
	15-19			20-24			25-29			30-34			35-39			40-44			45-49			Observed		Standardized ^b			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																											
Bangladesh	92	4	4	84	8	8	84	6	10	78	9	13	78	9	13	82	7	11	87	8	5	84	7	9	83	7	10
Fiji	59	10	31	37	10	53	25	10	65	21	11	68	27	10	63	31	13	56	46	14	40	31	11	58	31	11	58
Indonesia	82	3	15	62	3	35	50	4	46	51	5	44	57	4	39	66	5	29	79	6	15	62	4	34	59	4	36
Jordan	83	5	12	61	9	30	47	9	44	43	8	49	46	8	46	46	5	49	57	5	38	53	7	40	51	8	41
Korea, Rep. of	83	4	13	70	4	26	46	7	47	30	5	65	24	4	72	34	4	62	60	4	36	41	5	54	45	5	50
Malaysia	75	9	16	54	10	36	41	10	49	40	10	50	47	10	43	50	11	39	69	12	19	50	10	40	49	11	40
Nepal	99	0	1	98	1	1	95	1	4	93	1	6	92	1	7	94	1	5	97	0	3	95	1	4	95	1	4
Pakistan ^c	99	0	1	96	0	4	89	2	9	86	2	12	82	2	16	84	4	12	88	2	10	89	2	9	89	2	9
Philippines	74	13	13	48	16	36	34	19	47	33	17	50	33	19	48	44	20	36	61	18	21	42	18	40	41	18	41
Sri Lanka	80	9	11	67	12	21	57	17	26	43	21	36	44	20	35	54	21	25	63	24	13	54	19	27	55	18	27
Thailand	73	7	20	57	6	37	42	6	51	42	7	51	44	4	52	58	7	35	72	10	18	52	7	41	51	6	43
<i>Caribbean and Latin America</i>																											
Colombia	55	9	36	36	14	50	33	10	57	25	11	64	35	10	55	43	16	41	57	17	26	38	12	50	37	12	51
Costa Rica	NA	NA	NA	14	6	80	10	7	83	12	8	80	16	9	75	18	12	70	37	14	49	16	9	75	16	9	75
Dominican Rep.	70	12	18	51	14	35	39	9	52	41	11	48	46	9	45	56	10	34	69	9	22	50	11	39	49	11	40
Guyana	62	12	27	48	10	42	38	9	53	32	7	61	41	8	51	50	8	42	54	12	34	45	9	46	43	9	48
Jamaica	38	9	53	28	11	61	22	8	70	22	9	69	33	9	58	46	8	46	52	13	35	33	9	58	31	9	60
Mexico	74	7	19	55	9	36	43	11	46	44	10	46	50	11	39	60	10	30	71	10	19	53	10	37	52	10	38
Panama	NA	NA	NA	28	12	60	20	9	71	24	8	68	24	11	65	29	10	61	35	15	50	26	10	64	26	10	64
Peru	66	23	11	51	29	20	42	29	29	40	26	34	49	24	27	54	27	19	66	20	14	50	26	24	49	26	25

Table 1.2.2 Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Ever Used, by Number of Living Children

Country	Number of Living Children and Type of Method ^a Ever Used												Total														
	None			1			2			3			4			5+			Observed		Standardized ^b						
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E			
<i>Asia and Pacific</i>																											
Bangladesh	95	3	2	89	6	5	86	7	7	83	8	9	82	9	9	77	9	14	84	7	9	84	7	9	84	7	9
Fiji	74	9	17	41	14	45	28	12	60	24	10	66	21	9	70	18	11	71	31	11	58	31	11	58	31	11	58
Indonesia	94	1	5	71	4	25	55	4	41	50	5	45	53	6	41	51	5	44	62	4	34	59	4	36	59	4	36
Jordan	87	6	7	66	8	26	55	7	38	49	9	42	50	7	43	46	7	47	53	7	40	56	7	37	56	7	37
Korea, Rep. of	80	4	16	71	6	23	39	7	54	28	5	67	27	4	69	35	4	61	41	5	54	46	5	49	46	5	49
Malaysia	84	8	8	57	10	33	46	12	42	44	11	45	44	12	44	45	10	45	50	10	40	51	11	38	51	11	38
Nepal	99	0	1	98	1	1	97	1	2	94	1	5	92	1	7	89	1	10	95	1	4	94	1	5	94	1	5
Pakistan ^c	99	0	1	97	1	2	94	2	4	90	1	9	87	2	11	80	3	17	89	2	9	89	2	9	89	2	9
Philippines	88	5	7	51	20	29	40	20	40	31	20	49	33	18	49	40	18	42	42	18	40	45	17	38	45	17	38
Sri Lanka	92	6	2	67	19	14	51	26	23	46	23	31	42	20	38	48	17	35	54	19	27	55	18	26	55	18	26
Thailand	83	6	11	56	7	37	47	7	46	44	6	50	42	7	51	53	6	41	52	7	41	54	7	41	54	7	40
<i>Caribbean and Latin America</i>																											
Colombia	65	10	25	41	11	48	28	14	58	33	11	56	30	14	56	39	13	48	38	12	50	39	12	49	39	12	49
Costa Rica ^e	55	6	39	12	7	81	9	8	83	10	10	80	12	9	79	19	10	71	16	9	75	18	9	73	18	9	73
Dominican Rep.	77	8	15	62	10	28	39	14	47	37	13	50	39	10	51	49	10	41	50	11	39	49	11	40	49	11	40
Guyana	61	10	29	55	11	34	41	9	50	40	11	49	40	8	52	39	7	53	45	9	46	44	9	47	44	9	47
Jamaica	50	11	39	41	10	49	25	9	66	29	8	63	26	13	61	28	8	64	33	9	58	31	9	59	31	9	59
Mexico	80	6	14	56	9	35	43	11	46	47	9	44	49	12	39	54	11	35	53	10	37	54	10	36	54	10	36
Panama ^e	53	11	36	27	7	66	21	9	70	20	9	71	20	10	70	28	12	60	26	10	64	28	10	62	28	10	62
Peru	78	13	9	51	30	19	42	28	30	42	26	32	46	27	27	55	25	20	50	26	24	52	25	22	52	25	22

GROUP 2 EVER-USE BY INDIVIDUAL METHODS

LIST OF TABLES

2.1.1	Percentage of Currently Married Women Who Have Ever Used the Pill, by Current Age	31	2.7.1	Percentage of Currently Married Women Who Have Ever Used Withdrawal, by Current Age	37
2.1.2	Percentage of Currently Married Women Who Have Ever Used the Pill, by Number of Living Children	31	2.7.2	Percentage of Currently Married Women Who Have Ever Used Withdrawal, by Number of Living Children	37
2.2.1	Percentage of Currently Married Women Who Have Ever Used an IUD, by Current Age	32	2.8.1	Percentage of Currently Married Women Who Have Ever Used Abstinence, by Current Age	38
2.2.2	Percentage of Currently Married Women Who Have Ever Used an IUD, by Number of Living Children	32	2.8.2	Percentage of Currently Married Women Who Have Ever Used Abstinence, by Number of Living Children	38
2.3.1	Percentage of Currently Married Women Who Have Ever Used Other Female Scientific Methods, by Current Age	33	2.9.1	Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Current Age	39
2.3.2	Percentage of Currently Married Women Who Have Ever Used Other Female Scientific Methods, by Number of Living Children	33	2.9.2	Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Number of Living Children	39
2.4.1	Percentage of Currently Married Women Who Have Ever Used Douche, by Current Age	34	2.10.1	Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Current Age	40
2.4.2	Percentage of Currently Married Women Who Have Ever Used Douche, by Number of Living Children	34	2.10.2	Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Number of Living Children	40
2.5.1	Percentage of Currently Married Women Who Have Ever Used the Condom, by Current Age	35	2.11.1	Percentage of Currently Married Women Who Have Ever Used Injection, by Current Age	41
2.5.2	Percentage of Currently Married Women Who Have Ever Used the Condom, by Number of Living Children	35	2.11.2	Percentage of Currently Married Women Who Have Ever Used Injection, by Number of Living Children	41
2.6.1	Percentage of Currently Married Women Who Have Ever Used Rhythm, by Current Age	36	2.12.1	Percentage of Currently Married Women Who Have Ever Used Other Methods, by Current Age	42
2.6.2	Percentage of Currently Married Women Who Have Ever Used Rhythm, by Number of Living Children	36	2.12.2	Percentage of Currently Married Women Who Have Ever Used Other Methods, by Number of Living Children	42

Table 2.1.1 Percentage of Currently Married Women Who Have Ever Used the Pill, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	2.3	5.9	7.8	9.7	6.8	6.5	1.7	5.9
Fiji	17.9	36.5	40.2	35.4	31.1	21.9	12.8	31.6
Indonesia	13.0	28.4	35.8	33.1	28.1	20.4	11.3	26.2
Jordan	11.3	28.1	40.7	45.5	41.7	44.3	32.5	36.5
Korea, Rep. of	9.1	16.6	30.8	45.1	50.2	35.7	17.7	35.0
Malaysia	15.3	31.3	43.7	42.9	36.7	29.9	12.3	34.1
Nepal	0.3	1.1	2.3	2.6	3.7	3.1	1.4	2.0
Pakistan ^C	0.4	1.8	3.9	5.7	7.0	4.7	2.4	3.7
Philippines	4.3	19.7	31.6	30.9	30.9	22.4	12.8	25.2
Sri Lanka	1.9	7.8	10.6	13.2	9.9	5.8	3.1	8.7
Thailand	16.9	27.0	37.0	33.2	33.2	19.2	9.5	27.3
<i>Caribbean and Latin America</i>								
Colombia	27.6	37.6	42.6	44.4	34.2	22.8	14.2	34.5
Costa Rica	NA	68.2	68.8	59.6	49.0	32.3	17.3	53.0
Dominican Rep.	13.5	25.6	32.8	22.9	18.4	11.6	7.5	21.3
Guyana	13.8	25.3	31.1	29.2	18.8	11.4	7.9	21.8
Jamaica	26.5	32.2	45.8	41.8	28.6	18.4	11.8	31.1
Mexico	14.0	26.9	34.2	35.0	28.0	20.6	11.0	26.7
Panama	NA	51.4	59.6	50.9	38.9	25.1	12.5	44.0
Peru	6.5	13.6	18.6	20.4	13.6	10.7	5.2	13.9

Table 2.1.2 Percentage of Currently Married Women Who Have Ever Used the Pill, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	1.0	3.6	5.8	6.1	6.2	9.6	5.9
Fiji	11.1	29.4	36.8	41.6	36.9	31.4	31.6
Indonesia	5.0	21.8	33.3	34.2	29.7	30.6	26.2
Jordan	5.8	22.0	34.3	38.3	40.4	43.3	36.5
Korea, Rep. of	10.0	15.2	32.1	45.3	45.1	40.1	35.0
Malaysia	4.6	27.1	36.5	39.0	38.0	38.7	34.1
Nepal	0.1	0.6	0.9	2.9	3.7	5.1	2.0
Pakistan ^C	0.3	0.9	1.7	4.0	5.1	7.1	3.7
Philippines	4.5	14.2	23.9	31.2	30.9	27.6	25.2
Sri Lanka	1.0	4.4	7.9	11.0	13.9	10.2	8.7
Thailand	6.6	28.8	31.4	33.2	28.9	26.5	27.3
<i>Caribbean and Latin America</i>							
Colombia	20.1	36.1	42.3	38.6	39.3	30.3	34.5
Costa Rica ^e	30.8	66.2	66.0	57.8	55.4	40.5	53.0
Dominican Rep.	10.8	22.3	30.9	27.2	25.6	16.8	21.3
Guyana	14.1	22.3	29.5	27.0	24.6	18.8	21.8
Jamaica	19.4	25.4	39.6	40.3	32.9	30.3	31.1
Mexico	10.1	25.3	35.4	31.3	30.5	24.7	26.7
Panama ^e	27.7	54.8	54.7	51.2	48.2	32.3	44.0
Peru	5.0	12.0	19.6	18.9	15.6	10.3	13.9

Table 2.2.1 Percentage of Currently Married Women Who Have Ever Used an IUD, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.5	0.5	1.7	3.4	3.0	1.5	1.2
Fiji	0.9	8.4	19.3	25.3	18.9	16.8	10.6	16.5
Indonesia	1.5	6.3	9.9	11.2	10.9	7.7	4.5	8.0
Jordan	0.7	0.7	4.8	6.6	6.5	7.5	4.0	4.5
Korea, Rep. of	1.8	7.0	17.3	29.3	39.8	35.4	19.1	25.7
Malaysia	0.0	0.6	1.4	2.7	1.9	2.3	1.0	1.6
Nepal	0.0	0.0	0.1	0.6	0.6	0.3	0.5	0.2
Pakistan ^c	0.0	0.5	2.1	4.2	6.4	5.1	5.1	3.1
Philippines	1.8	4.3	7.6	9.4	9.1	7.8	3.9	7.2
Sri Lanka	5.4	8.4	7.9	10.9	10.8	6.7	2.7	8.1
Thailand	3.5	8.9	9.9	16.8	13.7	6.6	3.5	9.9
<i>Caribbean and Latin America</i>								
Colombia	5.5	14.0	21.0	22.1	18.7	13.8	6.9	16.3
Costa Rica	NA	10.8	14.4	15.9	15.9	11.4	6.2	12.9
Dominican Rep.	2.4	4.6	11.2	9.0	10.0	2.9	3.1	6.9
Guyana	1.5	8.3	11.0	14.7	10.9	4.5	5.9	8.8
Jamaica	1.2	4.0	7.5	14.6	14.3	7.6	8.4	8.1
Mexico	2.5	7.5	13.6	12.3	9.2	5.6	1.3	8.6
Panama	NA	7.5	12.2	10.4	11.2	5.7	2.9	9.1
Peru	0.4	2.0	2.9	4.9	3.2	2.4	0.7	2.7

Table 2.2.2 Percentage of Currently Married Women Who Have Ever Used an IUD, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.3	0.3	0.5	1.3	2.9	1.2
Fiji	0.0	5.5	14.3	19.4	27.2	22.9	16.5
Indonesia	0.1	3.5	7.2	11.4	11.6	12.8	8.0
Jordan	0.0	1.2	4.8	3.7	5.3	5.8	4.5
Korea, Rep. of	0.6	5.3	21.0	32.3	37.9	33.7	25.7
Malaysia	0.4	0.8	1.6	1.9	1.7	2.0	1.6
Nepal	0.0	0.0	0.2	0.0	0.3	1.1	0.2
Pakistan ^c	0.3	0.5	0.7	2.2	3.1	7.2	3.1
Philippines	0.2	3.2	7.0	9.1	9.5	7.8	7.2
Sri Lanka	0.1	5.1	9.4	9.4	11.4	9.3	8.1
Thailand	0.0	5.6	11.6	16.7	12.0	9.7	9.9
<i>Caribbean and Latin America</i>							
Colombia	0.0	10.3	22.3	22.2	20.1	16.2	16.3
Costa Rica ^e	1.6	9.8	13.3	14.8	12.6	15.8	12.9
Dominican Rep.	0.0	3.4	8.1	10.6	9.4	8.2	6.9
Guyana	0.2	2.9	9.1	10.2	12.5	12.9	8.8
Jamaica	1.0	1.4	6.1	9.7	11.0	15.1	8.1
Mexico	0.6	7.7	11.6	11.6	11.2	7.7	8.6
Panama ^e	0.6	6.5	11.5	9.9	11.9	8.8	9.1
Peru	0.0	2.2	4.6	2.5	3.5	2.1	2.7

Table 2.3.1 Percentage of Currently Married Women Who Have Ever Used Other Female Scientific Methods, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	1.0	1.0	0.5	0.3	0.8	0.1	0.6
Fiji ^f	0.4	2.1	3.7	3.9	3.6	5.5	3.5	3.5
Indonesia	0.0	0.2	0.2	0.6	0.7	0.1	0.1	0.3
Jordan	0.7	1.6	1.0	1.7	2.2	3.0	1.9	1.7
Korea, Rep. of	0.0	0.2	0.6	1.0	0.7	0.5	0.2	0.6
Malaysia	0.0	0.2	0.3	1.9	2.0	4.3	3.9	1.8
Nepal	NAsk							
Pakistan ^c	0.1	1.0	0.7	0.8	2.1	0.9	0.6	0.9
Philippines	0.0	1.1	3.0	4.3	4.0	2.8	2.2	3.0
Sri Lanka	0.0	0.0	0.2	0.4	0.2	0.5	1.0	0.3
Thailand	0.6	0.5	2.4	0.5	1.2	0.6	0.4	1.0
<i>Caribbean and Latin America</i>								
Colombia	7.2	9.1	14.7	18.9	14.8	12.4	3.6	12.5
Costa Rica	NA	11.0	12.6	17.1	11.2	10.3	6.8	12.0
Dominican Rep.	2.9	4.1	10.1	11.7	9.0	6.4	3.7	7.2
Guyana	5.5	6.5	9.2	10.7	12.1	9.5	8.2	8.8
Jamaica	4.9	6.9	14.7	16.2	13.0	11.9	10.1	11.2
Mexico	2.5	6.8	10.3	12.3	8.4	6.0	4.2	8.0
Panama	NA	9.1	11.3	10.4	11.7	10.1	4.0	9.9
Peru	2.6	3.3	5.6	7.4	5.8	4.5	3.6	5.0

Table 2.3.2 Percentage of Currently Married Women Who Have Ever Used Other Female Scientific Methods, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.3	0.6	0.8	0.7	0.9	0.6
Fiji ^f	1.3	1.4	4.5	4.3	3.2	4.4	3.5
Indonesia	0.0	0.0	0.1	0.2	0.9	0.7	0.3
Jordan	0.0	1.4	1.6	2.5	1.7	1.9	1.7
Korea, Rep. of	0.3	0.6	0.6	0.8	0.6	0.6	0.6
Malaysia	0.0	0.8	0.9	1.4	3.1	2.6	1.8
Nepal	NAsk						
Pakistan ^c	0.1	0.2	0.3	1.2	0.7	1.8	0.9
Philippines	0.0	0.9	3.1	4.3	4.0	3.1	3.0
Sri Lanka	0.0	0.0	0.3	0.2	1.0	0.4	0.3
Thailand	0.0	1.4	1.4	0.9	1.3	0.7	1.0
<i>Caribbean and Latin America</i>							
Colombia	6.8	10.1	18.6	14.0	14.9	10.6	12.5
Costa Rica ^e	6.6	10.7	15.6	17.0	13.3	8.7	12.0
Dominican Rep.	3.1	3.8	10.4	8.9	6.9	8.3	7.2
Guyana	7.3	5.5	8.8	8.3	11.9	10.1	8.8
Jamaica	5.6	6.5	11.2	15.1	11.9	14.6	11.2
Mexico	4.5	6.4	11.9	11.4	9.9	6.1	8.0
Panama ^e	6.4	12.8	11.1	13.3	7.8	8.0	9.9
Peru	3.8	3.6	7.0	7.3	4.4	3.9	5.0

Table 2.4.1 Percentage of Currently Married Women Who Have Ever Used Douche, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.7	2.9	3.1	4.0	3.7	1.8	0.9	2.7
Fiji	Included in Table 2,3,1							
Indonesia	0.7	0.7	0.4	0.5	0.3	0.1	0.4	0.5
Jordan	1.4	3.1	2.8	2.4	3.4	1.9	1.6	2.5
Korea, Rep. of	1.8	2.5	4.3	4.9	5.3	4.1	1.7	4.1
Malaysia	N Ask							
Nepal								
Pakistan								
Philippines	1.1	2.0	1.9	3.1	2.0	2.3	2.0	2.2
Sri Lanka	0.0	1.6	1.8	1.5	1.6	1.6	1.4	1.5
Thailand	1.1	1.2	4.0	3.2	1.8	1.8	2.3	2.4
<i>Caribbean and Latin America</i>								
Colombia	5.0	3.6	4.7	5.9	7.3	5.6	5.5	5.3
Costa Rica	NA	3.0	4.6	5.5	5.4	5.7	5.9	4.9
Dominican Rep.	4.1	2.0	3.7	4.5	3.0	3.5	3.7	3.4
Guyana	2.5	3.0	4.2	4.0	5.6	4.7	9.5	4.5
Jamaica	1.6	4.0	3.2	5.8	5.1	3.6	4.6	4.0
Mexico	1.1	2.4	3.4	5.1	6.0	5.7	6.6	4.3
Panama	NA	8.5	9.5	9.2	11.5	13.4	11.7	10.3
Peru	9.6	13.9	16.6	18.6	18.2	13.9	10.1	15.2

Table 2.4.2 Percentage of Currently Married Women Who Have Ever Used Douche, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.7	2.5	2.6	3.3	3.4	3.2	2.7
Fiji	Included in Table 2.3.2						
Indonesia	0.4	0.3	0.5	0.5	0.9	0.3	0.5
Jordan	0.8	2.9	3.6	2.1	4.0	2.3	2.5
Korea, Rep. of	1.7	2.3	4.7	6.9	4.5	2.8	4.1
Malaysia	N Ask						
Nepal							
Pakistan							
Philippines	0.7	2.3	2.9	2.1	3.2	1.8	2.2
Sri Lanka	0.2	1.7	2.2	1.8	1.7	1.3	1.5
Thailand	1.2	3.1	4.0	2.1	2.5	1.6	2.4
<i>Caribbean and Latin America</i>							
Colombia	2.4	2.7	5.8	7.0	6.8	5.7	5.3
Costa Rica ^e	1.1	3.3	5.4	6.4	7.0	4.7	4.9
Dominican Rep.	4.0	1.5	3.9	4.9	4.4	2.8	3.4
Guyana	2.6	4.6	4.1	5.1	4.8	5.1	4.5
Jamaica	2.0	4.8	5.1	3.7	3.2	4.3	4.0
Mexico	2.5	2.0	4.6	5.6	4.6	4.9	4.3
Panama ^e	4.6	11.3	11.8	10.8	10.2	10.0	10.3
Peru	5.7	12.9	17.9	18.4	18.1	13.8	15.2

Table 2.5.1 Percentage of Currently Married Women Who Have Ever Used the Condom, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.3	2.8	2.6	2.7	2.7	2.2	1.4	2.3
Fiji	17.5	23.2	22.7	17.9	12.2	8.7	3.5	16.7
Indonesia	2.3	4.5	7.6	6.9	6.0	5.7	2.2	5.4
Jordan	3.0	5.7	9.6	11.3	10.9	9.9	10.6	8.9
Korea, Rep. of	3.6	9.9	23.2	29.5	29.2	21.3	12.1	22.5
Malaysia	2.1	7.7	11.0	12.0	9.0	10.0	4.4	9.1
Nepal	0.8	0.6	0.8	0.9	0.5	0.4	0.0	0.7
Pakistan ^c	0.1	1.6	4.3	3.7	4.3	2.3	0.6	2.6
Philippines	9.0	21.1	25.8	25.5	24.3	16.6	7.3	20.7
Sri Lanka	3.2	5.1	6.6	8.1	5.3	3.2	0.8	5.1
Thailand	2.6	4.4	4.8	4.4	4.3	2.5	0.3	3.6
<i>Caribbean and Latin America</i>								
Colombia	4.4	6.6	10.0	10.4	11.9	7.3	7.3	8.8
Costa Rica	NA	37.6	43.0	41.2	35.3	34.3	23.1	37.0
Dominican Rep.	5.3	8.4	13.5	16.5	9.4	8.7	6.8	10.3
Guyana	12.0	18.2	19.9	19.3	18.1	11.4	9.9	16.5
Jamaica	33.9	36.3	30.3	27.7	20.0	12.3	7.6	25.6
Mexico	1.4	4.2	7.1	9.4	8.3	7.2	6.1	6.6
Panama	NA	13.8	14.8	14.4	11.5	13.4	9.9	13.3
Peru	1.6	5.7	6.3	10.9	8.4	6.6	5.7	7.0

Table 2.5.2 Percentage of Currently Married Women Who Have Ever Used the Condom, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.8	1.4	2.5	2.9	2.7	3.0	2.3
Fiji	7.5	17.9	25.0	22.3	18.6	12.7	16.7
Indonesia	0.3	3.3	4.3	8.1	7.4	8.1	5.4
Jordan	1.3	7.6	8.7	14.1	11.0	8.9	8.9
Korea, Rep. of	6.4	11.4	24.8	31.1	27.3	21.7	22.5
Malaysia	4.3	8.5	11.6	11.3	10.5	7.9	9.1
Nepal	0.3	0.4	0.8	0.6	0.7	1.3	0.7
Pakistan ^c	0.2	0.7	2.5	2.7	3.9	4.2	2.6
Philippines	2.2	17.2	21.8	26.9	23.9	20.4	20.7
Sri Lanka	1.5	5.3	7.4	6.8	7.0	3.5	5.1
Thailand	4.1	5.5	4.1	4.8	3.3	1.7	3.6
<i>Caribbean and Latin America</i>							
Colombia	4.8	7.4	10.1	11.1	8.4	8.8	8.8
Costa Rica ^e	19.2	38.1	48.1	45.4	39.6	28.7	37.0
Dominican Rep.	4.0	6.8	15.8	13.8	15.8	8.7	10.3
Guyana	16.6	14.3	19.0	20.9	17.8	14.4	16.5
Jamaica	24.7	29.3	37.0	24.8	25.1	17.9	25.6
Mexico	2.3	3.6	8.9	9.6	7.7	6.3	6.6
Panama ^e	9.2	15.8	19.2	15.7	13.5	8.9	13.3
Peru	3.0	5.4	8.8	10.2	9.0	5.2	7.0

Table 2.6.1 Percentage of Currently Married Women Who Have Ever Used Rhythm, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	2.9	5.9	6.1	7.5	5.7	6.2	4.6	5.5
Fiji	6.7	15.2	17.5	18.4	16.7	16.2	9.6	15.8
Indonesia	0.2	2.5	4.3	4.0	4.2	2.6	1.8	3.0
Jordan	4.7	11.8	16.4	15.7	16.6	11.5	8.8	13.2
Korea, Rep. of	7.3	7.0	17.6	21.9	21.5	16.9	7.2	16.8
Malaysia	5.0	9.6	13.1	15.5	11.0	10.0	4.9	11.0
Nepal	N Ask							
Pakistan ^c	0.0	0.0	0.3	0.2	0.4	0.1	0.2	0.2
Philippines	6.8	17.7	27.6	28.1	29.0	20.6	16.2	23.5
Sri Lanka	6.2	14.1	20.3	29.1	27.8	23.2	21.1	22.5
Thailand	3.0	6.3	9.1	9.4	8.2	5.1	6.1	7.3
<i>Caribbean and Latin America</i>								
Colombia	7.7	21.1	21.5	23.1	20.1	19.2	14.9	19.7
Costa Rica	NA	15.4	21.9	22.2	23.5	26.6	22.5	21.7
Dominican Rep.	5.3	7.2	8.0	8.6	7.7	7.5	5.0	7.3
Guyana	8.9	11.2	9.7	15.3	10.7	11.4	9.5	11.1
Jamaica	4.1	5.8	5.7	6.4	2.5	5.8	5.1	5.2
Mexico	5.5	11.3	17.3	14.0	17.3	12.6	7.8	13.3
Panama	NA	16.2	17.1	16.8	15.9	15.5	16.1	16.4
Peru	14.4	27.8	34.7	33.9	26.9	24.7	15.8	27.4

Table 2.6.2 Percentage of Currently Married Women Who Have Ever Used Rhythm, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	1.9	4.0	5.8	7.2	6.8	6.4	5.5
Fiji	8.9	14.1	16.9	19.7	19.5	15.5	15.8
Indonesia	0.4	1.4	3.7	3.6	5.2	3.9	3.0
Jordan	4.3	8.0	18.3	17.6	19.0	12.3	13.2
Korea, Rep. of	9.7	9.1	19.9	24.3	18.3	13.9	16.8
Malaysia	7.6	12.5	14.9	14.7	11.5	7.9	11.0
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.3	0.2	0.5	0.1	0.2
Philippines	5.2	20.3	24.8	29.5	28.1	22.7	23.5
Sri Lanka	5.6	19.3	27.9	28.7	27.3	21.7	22.5
Thailand	4.4	7.9	9.1	8.6	9.2	5.3	7.3
<i>Caribbean and Latin America</i>							
Colombia	11.6	20.1	23.1	22.5	24.4	17.1	19.7
Costa Rica ^e	7.1	21.2	25.1	27.9	28.1	18.0	21.7
Dominican Rep.	5.8	6.0	9.7	8.9	8.9	6.2	7.3
Guyana	11.5	11.3	12.0	11.7	10.5	10.5	11.1
Jamaica	9.5	6.7	3.7	7.0	5.0	2.2	5.2
Mexico	6.8	10.9	17.4	17.0	15.8	11.9	13.3
Panama ^e	12.7	18.2	17.7	20.2	17.0	13.5	16.4
Peru	12.2	32.1	33.6	31.0	27.5	23.4	27.4

Table 2.7.1 Percentage of Currently Married Women Who Have Ever Used Withdrawal, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	2.0	3.6	3.0	3.5	3.4	3.1	1.6	3.0
Fiji	11.7	18.7	18.1	16.3	14.7	12.8	7.8	15.5
Indonesia	1.6	2.4	2.9	2.5	2.0	1.8	0.8	2.1
Jordan	7.4	11.7	20.7	25.9	23.1	20.8	16.9	18.9
Korea, Rep. of	5.5	4.9	12.6	15.9	16.1	12.1	6.6	12.3
Malaysia	5.0	7.8	9.5	9.8	7.7	9.8	4.4	8.3
Nepal	N Ask							
Pakistan ^c	0.0	0.3	0.2	0.4	0.4	0.4	0.0	0.3
Philippines	12.8	28.4	36.3	37.6	35.9	30.7	18.8	31.7
Sri Lanka	2.1	5.7	7.2	8.8	7.9	4.8	4.3	6.5
Thailand	5.2	4.2	6.8	5.5	5.0	5.4	5.2	5.4
<i>Caribbean and Latin America</i>								
Colombia	8.8	16.7	17.6	24.4	17.8	18.3	17.1	18.1
Costa Rica	NA	27.2	28.8	27.5	27.1	27.1	20.2	26.8
Dominican Rep.	11.8	18.2	23.6	23.7	20.4	16.8	11.2	19.0
Guyana	12.3	11.3	16.0	17.3	14.4	13.6	11.2	13.9
Jamaica	19.6	27.3	22.6	23.2	16.8	14.4	13.1	20.6
Mexico	10.8	16.2	19.8	22.2	21.1	19.1	12.9	18.3
Panama	NA	21.1	20.7	17.8	17.3	17.9	13.2	18.5
Peru	14.3	16.9	20.0	20.2	16.7	13.5	12.0	16.8

Table 2.7.2 Percentage of Currently Married Women Who Have Ever Used Withdrawal, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	1.1	2.5	3.4	3.3	3.8	3.3	3.0
Fiji	8.4	15.1	20.6	17.7	17.5	14.3	15.5
Indonesia	0.3	1.4	2.4	3.5	3.1	2.3	2.1
Jordan	6.4	13.1	17.1	20.3	19.7	21.8	18.9
Korea, Rep. of	5.3	7.0	14.6	17.1	14.2	10.6	12.3
Malaysia	5.0	6.3	10.9	9.2	9.9	7.7	8.3
Nepal	N Ask						
Pakistan ^c	0.0	0.1	0.8	0.2	0.4	0.2	0.3
Philippines	4.9	23.5	33.1	39.6	36.7	32.6	31.7
Sri Lanka	2.3	7.1	9.5	7.9	6.3	5.6	6.5
Thailand	3.5	5.6	5.7	6.5	8.0	4.0	5.4
<i>Caribbean and Latin America</i>							
Colombia	4.8	14.3	20.2	21.7	22.4	19.1	18.1
Costa Rica ^e	9.9	27.1	29.9	27.7	28.8	27.1	26.8
Dominican Rep.	6.3	15.5	24.7	23.6	23.6	19.4	19.0
Guyana	11.2	12.0	12.9	16.1	15.0	15.1	13.9
Jamaica	18.4	17.5	23.9	19.8	27.4	19.7	20.6
Mexico	5.8	14.9	20.5	18.7	22.7	19.9	18.3
Panama ^e	12.1	19.0	19.7	18.5	20.8	18.0	18.5
Peru	6.1	16.9	19.3	19.7	20.4	14.7	16.8

Table 2.8.1 Percentage of Currently Married Women Who Have Ever Used Abstinence, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	2.8	5.5	4.7	5.8	7.2	6.8	5.4	5.1
Fiji	1.8	7.5	9.0	11.9	11.4	13.7	9.3	9.9
Indonesia	1.4	2.7	3.8	4.4	4.8	4.5	4.1	3.7
Jordan	0.1	2.8	3.9	4.0	6.0	4.8	3.9	3.8
Korea, Rep. of	0.0	1.4	1.9	1.6	4.1	2.9	3.0	2.4
Malaysia	0.8	2.7	3.3	4.1	5.4	7.1	7.3	4.5
Nepal	0.0	0.4	0.0	0.6	0.5	0.7	0.0	0.3
Pakistan ^c	0.2	0.3	1.5	2.1	2.3	3.6	2.2	1.7
Philippines	2.1	7.6	10.5	11.5	12.8	11.1	9.0	10.4
Sri Lanka	3.3	6.7	9.2	13.6	14.0	14.2	17.1	12.1
Thailand	1.5	6.5	8.7	8.0	7.6	6.9	6.0	7.1
<i>Caribbean and Latin America</i>								
Colombia	4.4	4.7	5.9	7.4	5.9	6.8	5.1	5.9
Costa Rica	NA	2.6	4.7	3.7	4.5	4.9	4.6	4.1
Dominican Rep.	N Ask							
Guyana	5.2	8.9	7.4	8.5	7.0	7.0	6.6	7.5
Jamaica	9.4	14.6	9.5	10.4	12.1	7.2	8.9	10.7
Mexico	N Ask							
Panama	NA	9.9	8.5	7.9	4.9	6.9	7.0	7.7
Peru	2.5	6.4	7.2	7.8	8.2	9.0	8.4	7.5

Table 2.8.2 Percentage of Currently Married Women Who Have Ever Used Abstinence, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	1.2	3.0	6.3	5.8	6.5	6.7	5.1
Fiji	0.9	5.7	11.1	12.0	12.4	12.6	9.9
Indonesia	0.4	2.8	3.8	5.0	5.5	4.6	3.7
Jordan	1.7	2.8	2.1	3.4	4.8	4.6	3.8
Korea, Rep. of	1.7	1.3	1.9	1.9	3.2	3.6	2.4
Malaysia	0.7	2.5	3.3	5.3	4.8	6.2	4.5
Nepal	0.1	0.5	0.2	0.3	0.4	0.6	0.3
Pakistan ^c	0.0	0.6	1.4	1.3	2.4	3.0	1.7
Philippines	2.4	7.3	9.6	12.7	11.3	11.5	10.4
Sri Lanka	1.8	8.0	12.4	14.2	13.8	15.3	12.1
Thailand	2.5	5.9	9.1	7.9	8.2	6.9	7.1
<i>Caribbean and Latin America</i>							
Colombia	1.6	5.2	6.2	5.3	9.1	6.3	5.9
Costa Rica ^e	1.6	4.8	4.4	3.2	3.5	4.7	4.1
Dominican Rep.	N Ask						
Guyana	8.9	6.5	7.0	8.3	5.4	7.9	7.5
Jamaica	8.9	10.8	12.8	12.4	9.1	10.0	10.7
Mexico	N Ask						
Panama ^e	2.3	7.4	9.2	7.5	8.1	8.0	7.7
Peru	1.9	5.1	5.6	6.6	10.9	9.4	7.5

Table 2.9.1 Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.1	0.3	0.6	0.9	0.6	0.2	0.3
Fiji	0.0	0.9	9.9	22.1	27.8	26.6	20.7	15.8
Indonesia	0.0	0.0	0.1	0.3	0.7	0.4	0.3	0.3
Jordan	0.0	0.0	0.4	1.6	3.6	4.6	3.5	1.8
Korea, Rep. of	0.0	0.0	0.3	3.3	2.5	2.0	1.9	1.7
Malaysia	0.0	0.3	1.2	4.9	6.1	5.7	3.5	3.4
Nepal	0.0	0.1	0.0	0.0	0.4	0.0	0.0	0.1
Pakistan ^c	0.0	0.0	0.4	0.7	2.0	1.7	3.0	0.9
Philippines	0.0	1.4	4.4	6.3	7.2	5.3	2.7	4.7
Sri Lanka	0.5	1.7	7.7	13.1	14.5	11.6	6.2	9.2
Thailand	0.0	1.5	6.4	8.6	9.7	8.1	4.7	6.1
<i>Caribbean and Latin America</i>								
Colombia	0.0	0.8	2.2	6.6	8.2	6.2	2.9	4.0
Costa Rica	NA	0.6	6.8	13.1	23.0	20.3	15.3	12.3
Dominican Rep.	0.0	3.8	13.5	19.9	21.7	12.7	8.7	11.9
Guyana	0.0	0.3	3.0	12.9	17.0	19.8	14.5	8.5
Jamaica	0.0	0.6	2.5	13.4	16.2	19.1	10.1	8.1
Mexico	0.2	0.5	1.3	3.6	5.0	5.3	3.4	2.7
Panama	NA	0.8	13.6	23.8	31.4	35.8	35.5	21.2
Peru	0.0	0.2	1.3	4.2	6.3	2.9	2.9	2.8

Table 2.9.2 Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.0	0.0	0.3	0.5	0.6	0.3
Fiji	0.4	0.5	3.4	8.0	22.7	34.3	15.8
Indonesia	0.0	0.0	0.1	0.1	0.4	0.8	0.3
Jordan	0.5	0.0	0.4	0.6	0.5	3.1	1.8
Korea, Rep. of	0.0	0.1	2.4	2.5	2.2	1.7	1.7
Malaysia	0.0	0.4	1.0	1.6	4.4	6.7	3.4
Nepal	0.0	0.0	0.1	0.1	0.0	0.2	0.1
Pakistan ^c	0.1	0.1	0.6	0.8	0.9	1.9	0.9
Philippines	0.4	0.8	3.0	6.8	7.2	5.3	4.7
Sri Lanka	0.0	0.4	2.3	9.3	14.0	17.4	9.2
Thailand	0.0	0.4	3.8	8.4	12.0	8.7	6.1
<i>Caribbean and Latin America</i>							
Colombia	0.8	0.0	1.4	5.3	5.2	6.8	4.0
Costa Rica ^e	1.6	1.4	3.7	13.8	18.2	22.2	12.3
Dominican Rep.	0.9	0.8	8.1	17.5	19.7	17.6	11.9
Guyana	0.5	0.4	1.8	5.6	7.1	19.2	8.5
Jamaica	0.7	1.7	1.9	4.7	8.7	20.3	8.1
Mexico	0.4	0.6	1.4	2.6	2.7	4.5	2.7
Panama ^e	1.7	3.0	10.5	25.6	28.8	31.8	21.2
Peru	0.0	0.2	0.9	4.4	3.2	4.3	2.8

Table 2.10.1 Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.3	0.3	0.4	2.3	0.7	0.4	0.5
Fiji	0.0	0.0	0.0	0.3	0.0	0.2	0.0	0.1
Indonesia	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Jordan	0.0	0.0	0.0	0.0	0.1	0.1	0.9	0.1
Korea, Rep. of	0.0	0.2	1.4	2.8	6.0	5.5	4.0	3.3
Malaysia	0.0	0.0	0.2	0.2	0.7	0.8	0.7	0.4
Nepal	0.0	0.3	1.6	3.2	3.3	2.1	1.7	1.6
Pakistan ^c	0.0	0.0	0.0	0.1	0.2	0.2	0.0	0.1
Philippines	0.0	0.3	0.6	0.9	1.1	0.5	0.4	0.7
Sri Lanka	0.0	0.2	0.7	1.3	1.0	0.9	0.0	0.7
Thailand	0.0	0.9	1.9	1.6	4.9	2.6	1.4	2.1
<i>Caribbean and Latin America</i>								
Colombia	0.0	0.2	0.3	0.2	0.0	0.3	0.0	0.2
Costa Rica	NA	0.0	1.4	2.0	0.7	0.6	1.0	1.0
Dominican Rep.	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.1
Guyana	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.0	0.0	0.1	0.3	0.7	0.2	0.0	0.2
Panama	NA	0.2	0.0	0.9	0.5	0.3	0.4	0.4
Peru	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 2.10.2 Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.1	0.3	0.5	0.8	0.9	0.5
Fiji	0.0	0.0	0.1	0.3	0.2	0.0	0.1
Indonesia	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Jordan	0.0	0.0	0.0	0.1	0.0	0.2	0.1
Korea, Rep. of	0.0	0.3	2.4	4.6	5.3	4.1	3.3
Malaysia	0.0	0.1	0.1	0.3	0.3	0.7	0.4
Nepal	0.1	0.3	1.0	1.8	3.0	4.5	1.6
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.2	0.1
Philippines	0.0	0.4	0.3	0.6	1.0	0.8	0.7
Sri Lanka	0.0	0.1	0.6	1.0	1.5	0.8	0.7
Thailand	0.8	0.2	2.3	4.8	4.7	1.0	2.1
<i>Caribbean and Latin America</i>							
Colombia	0.0	0.0	0.0	0.2	0.3	0.3	0.2
Costa Rica ^e	0.0	0.7	0.6	1.7	1.1	1.1	1.0
Dominican Rep.	0.0	0.0	0.0	0.0	0.0	0.2	0.1
Guyana	0.0	0.0	0.2	0.0	0.0	0.0	0.0
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.2	0.1	0.4	0.1	0.1	0.2	0.2
Panama ^e	0.0	0.0	0.6	0.4	0.3	0.4	0.4
Peru	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 2.11.1 Percentage of Currently Married Women Who Have Ever Used Injection, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	N Ask							
Fiji	0.4	0.9	0.9	0.7	0.6	0.4	0.0	0.6
Indonesia	0.6	0.6	1.4	0.8	0.8	0.9	0.4	0.8
Jordan	N Ask							
Korea, Rep. of	0.0	0.2	0.5	0.9	0.6	0.5	0.2	0.5
Malaysia	0.8	0.9	1.3	1.6	1.3	1.1	0.7	1.2
Nepal	N Ask							
Pakistan ^c	0.0	0.2	0.4	0.3	0.7	0.5	0.6	0.4
Philippines	N Ask							
Sri Lanka	0.0	1.0	0.9	1.2	0.6	0.6	0.5	0.8
Thailand	1.9	5.1	7.4	4.2	5.5	3.3	2.5	4.7
<i>Caribbean and Latin America</i>								
Colombia	2.8	2.1	4.4	5.5	5.4	2.5	0.7	3.6
Costa Rica	NA	5.8	9.5	12.5	9.4	11.1	5.9	9.2
Dominican Rep.	1.2	1.5	1.7	0.4	0.0	0.0	0.6	0.9
Guyana	0.0	1.4	1.4	3.0	1.9	0.6	0.7	1.4
Jamaica	6.5	14.8	17.9	16.8	9.8	3.6	1.3	11.3
Mexico	3.2	6.5	11.1	10.4	8.3	5.0	2.3	7.5
Panama	NA	4.7	4.6	3.4	2.3	1.2	0.4	3.2
Peru	2.6	4.7	7.6	7.4	5.7	1.6	1.1	4.9

Table 2.11.2 Percentage of Currently Married Women Who Have Ever Used Injection, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	N Ask						
Fiji	0.0	0.8	1.0	0.8	1.3	0.4	0.6
Indonesia	0.0	0.5	0.9	1.2	0.9	1.3	0.8
Jordan	N Ask						
Korea, Rep. of	0.3	0.6	0.5	0.7	0.6	0.5	0.5
Malaysia	0.2	0.4	1.0	1.4	1.7	1.5	1.2
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.0	0.0	0.6	0.9	0.4
Philippines	N Ask						
Sri Lanka	0.2	0.4	1.3	1.2	0.6	0.7	0.8
Thailand	1.1	6.6	6.4	4.3	5.4	3.8	4.7
<i>Caribbean and Latin America</i>							
Colombia	1.2	2.0	4.5	2.2	5.5	4.5	3.6
Costa Rica ^e	1.1	6.0	8.9	11.4	10.5	11.1	9.2
Dominican Rep.	0.0	1.5	1.9	0.4	1.5	0.5	0.9
Guyana	0.5	0.4	1.6	1.9	1.1	2.0	1.4
Jamaica	1.6	6.5	18.1	11.7	16.0	13.2	11.3
Mexico	2.5	7.0	9.7	10.3	6.3	7.4	7.5
Panama ^e	3.5	3.9	5.6	3.7	2.2	1.8	3.2
Peru	0.4	3.6	5.3	7.8	6.1	4.2	4.9

Table 2.12.1 Percentage of Currently Married Women Who Have Ever Used Other Methods, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.2	0.8	2.2	1.1	1.0	0.7	0.7
Fiji	1.8	3.4	6.9	9.5	8.9	9.4	10.8	7.5
Indonesia	3.6	5.3	6.9	6.6	5.9	5.2	3.5	5.5
Jordan	1.4	6.3	9.1	14.2	13.1	14.8	10.5	10.2
Korea, Rep. of	0.0	0.2	0.2	0.2	0.7	1.1	1.3	0.5
Malaysia	3.3	4.2	5.2	4.8	6.0	6.0	4.7	5.1
Nepal	0.4	0.6	1.2	0.2	0.2	0.5	0.3	0.5
Pakistan ^c	0.0	0.1	0.3	0.4	0.5	0.7	0.5	0.3
Philippines	0.5	1.1	1.1	1.2	0.7	1.0	0.9	1.0
Sri Lanka	0.0	0.5	0.1	0.2	0.3	0.4	0.1	0.2
Thailand	1.2	0.6	0.9	1.3	0.7	0.6	0.3	0.8
<i>Caribbean and Latin America</i>								
Colombia	1.7	3.4	3.7	4.9	5.2	2.8	3.6	3.8
Costa Rica	NA	2.0	2.3	2.0	1.8	2.0	1.3	1.9
Dominican Rep.	0.6	4.9	9.2	9.8	11.7	10.4	6.8	7.9
Guyana	3.4	2.4	3.4	3.2	3.0	1.4	2.3	2.8
Jamaica	1.2	2.7	4.5	3.7	1.0	1.8	0.8	2.5
Mexico	0.7	0.4	0.7	0.9	0.8	0.3	0.6	0.6
Panama	NA	2.2	2.6	2.1	1.2	1.2	1.8	1.9
Peru	1.4	2.9	2.4	4.1	2.4	2.9	2.1	2.8

Table 2.12.2 Percentage of Currently Married Women Who Have Ever Used Other Methods, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.1	0.2	0.9	0.8	1.6	0.7
Fiji	0.0	4.1	5.5	7.1	10.3	11.7	7.5
Indonesia	0.8	4.6	5.3	7.2	6.9	7.5	5.5
Jordan	0.0	4.2	5.9	9.7	8.2	14.4	10.2
Korea, Rep. of	0.0	0.4	0.1	0.5	0.6	1.1	0.5
Malaysia	0.9	3.4	4.8	5.3	5.8	6.4	5.1
Nepal	0.3	0.5	0.4	0.8	1.0	0.5	0.5
Pakistan ^c	0.0	0.2	0.6	0.2	0.2	0.7	0.3
Philippines	0.0	0.7	1.3	1.5	1.3	0.8	1.0
Sri Lanka	0.2	0.2	0.1	0.3	0.3	0.3	0.2
Thailand	0.0	0.8	0.7	1.2	1.1	0.8	0.8
<i>Caribbean and Latin America</i>							
Colombia	0.8	2.0	4.7	3.6	4.5	4.8	3.8
Costa Rica ^e	0.0	1.4	1.5	1.7	3.2	2.5	1.9
Dominican Rep.	0.9	1.5	7.7	9.8	12.8	10.8	7.9
Guyana	4.0	1.9	1.6	3.2	2.8	3.1	2.8
Jamaica	1.6	3.1	4.0	2.0	3.7	1.3	2.5
Mexico	0.4	0.0	1.0	0.7	0.4	0.8	0.6
Panama ^e	2.3	0.6	1.9	1.7	1.9	2.5	1.9
Peru	0.5	1.4	3.0	2.4	3.1	3.6	2.8

GROUP 3 CURRENT USE BY TYPE OF METHOD

LIST OF TABLES

3.1.1	Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Currently Used, by Current Age	45
3.1.2	Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Currently Used, by Number of Living Children	45
3.2.1	Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Currently Used, by Current Age	46
3.2.2	Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Currently Used, by Number of Living Children	46
3.3.1	Per Cent Distribution of Currently Married Fecund Women according to Type of Contraceptive Method Currently Used, by Current Age	47
3.3.2	Per Cent Distribution of Currently Married Fecund Women according to Type of Contraceptive Method Currently Used, by Number of Living Children	47
3.4.1	Per Cent Distribution of Exposed Women according to Type of Contraceptive Method Currently Used, by Current Age	48
3.4.2	Per Cent Distribution of Exposed Women according to Type of Contraceptive Method Currently Used, by Number of Living Children	48

Table 3.1.1 Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Currently Used, by Current Age

Country	Current Age and Type of Method ^a Currently Used															Total											
	15-19			20-24			25-29			30-34			35-39			40-44			45-49			Observed		Standardized ^b			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																											
Bangladesh	96	2	2	93	3	4	92	2	6	89	4	7	89	4	7	93	3	4	96	2	2	93	3	4	92	3	5
Fiji	79	4	17	69	6	25	61	5	34	52	7	41	52	6	42	58	5	37	73	2	25	61	5	34	61	5	34
Indonesia	89	2	9	76	2	22	70	4	26	70	4	26	73	4	23	81	3	16	91	2	7	77	3	20	76	3	21
Jordan	92	2	6	84	5	11	74	7	19	69	11	20	68	10	22	70	9	21	83	6	11	76	7	17	75	8	17
Korea, Rep. of	87	4	9	87	3	10	72	6	22	56	10	34	49	11	40	67	9	24	89	1	10	68	7	25	70	7	23
Malaysia	86	3	11	72	7	21	63	8	29	63	10	27	65	10	25	70	11	19	87	6	7	69	9	22	68	9	23
Nepal	100	0	0	99	0	1	98	0	2	96	0	4	96	0	4	98	0	2	99	0	1	98	0	2	98	0	2
Pakistan ^c	100	0	0	98	0	2	95	1	4	94	2	4	91	2	7	93	3	4	94	2	4	95	1	4	95	1	4
Philippines	84	11	5	72	15	13	62	21	17	58	22	20	56	24	20	66	21	13	84	10	6	66	19	15	66	19	15
Sri Lanka	87	3	10	81	7	12	71	11	18	58	16	26	63	13	24	70	13	17	84	10	6	71	12	17	71	11	18
Thailand	83	2	15	71	3	26	61	3	36	60	3	37	62	2	36	73	3	24	89	2	9	69	3	28	68	3	29
<i>Caribbean and Latin America</i>																											
Colombia	78	5	17	64	10	26	58	10	32	50	13	37	59	9	32	67	14	19	83	8	9	63	10	27	62	11	28
Costa Rica	NA	NA	NA	41	9	50	38	10	52	37	9	54	36	11	53	43	13	44	64	7	29	42	10	48	41	10	49
Dominican Rep.	90	4	6	78	4	18	65	6	29	65	5	30	64	5	31	75	6	19	83	2	15	73	5	22	71	5	24
Guyana	84	7	9	77	7	16	69	6	25	60	8	32	63	9	28	68	5	27	78	4	18	71	6	23	69	7	24
Jamaica	75	1	24	67	2	31	62	2	36	55	3	42	58	3	39	68	2	30	75	1	24	65	2	33	64	2	34
Mexico	87	3	10	75	5	20	64	7	29	65	7	28	66	10	24	78	7	15	90	3	7	73	6	21	71	7	22
Panama	NA	NA	NA	62	8	30	48	7	45	49	6	45	47	7	46	48	7	45	57	6	37	52	7	41	52	7	41
Peru	85	10	5	73	19	8	66	22	12	62	22	16	68	19	13	74	19	7	86	10	4	72	18	10	71	19	11

Table 3.1.2 Per Cent Distribution of Ever-Married Women according to Type of Contraceptive Method Currently Used, by Number of Living Children

Country	Number of Living Children and Type of Method ^a Currently Used															Total								
	None			1			2			3			4			5+			Observed		Standardized ^d			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																								
Bangladesh	99	0	1	95	2	3	94	2	4	92	3	5	92	3	5	89	4	7	93	3	4	92	3	5
Fiji	95	2	3	74	6	20	64	8	28	60	6	34	49	7	44	46	5	49	61	5	34	61	5	34
Indonesia	98	0	2	83	2	15	73	3	24	69	4	27	71	4	25	69	4	27	77	3	20	75	3	22
Jordan	96	2	2	84	5	11	74	7	19	76	8	16	76	8	16	71	9	20	76	7	17	78	7	15
Korea, Rep. of	93	3	4	89	3	8	63	9	28	56	10	34	56	9	35	67	7	26	68	7	25	72	6	22
Malaysia	94	3	3	76	5	19	68	9	23	64	11	25	63	11	26	66	9	25	69	9	22	71	8	21
Nepal	100	0	0	99	0	1	98	0	2	98	0	2	97	0	3	94	0	6	98	0	2	97	0	3
Pakistan ^c	100	0	0	99	0	1	97	1	2	95	1	4	94	2	4	90	3	7	95	1	4	95	1	4
Philippines	97	2	1	76	14	10	62	21	17	56	24	20	58	21	21	65	20	15	66	19	15	68	18	14
Sri Lanka	98	1	1	81	11	8	70	16	14	64	15	21	60	15	25	66	10	24	71	12	17	72	11	17
Thailand	95	1	4	73	3	24	65	2	33	59	3	38	60	3	37	71	2	27	69	3	28	71	2	27
<i>Caribbean and Latin America</i>																								
Colombia	88	3	9	68	9	23	57	13	30	58	9	33	55	12	33	63	11	26	63	10	27	64	10	26
Costa Rica ^e	81	3	16	45	9	46	34	12	54	36	10	54	37	10	53	42	9	49	42	10	48	44	9	47
Dominican Rep.	92	2	6	85	3	12	71	4	25	63	7	30	64	6	30	68	5	27	73	5	22	72	5	23
Guyana	85	7	8	82	4	14	72	7	21	68	9	23	68	8	24	61	6	33	71	6	23	71	6	23
Jamaica	77	2	21	74	1	25	62	1	37	65	2	33	61	3	36	58	2	40	65	2	33	64	2	34
Mexico	94	1	5	78	5	17	65	7	28	66	7	27	67	8	25	73	7	20	73	6	21	73	6	21
Panama ^e	84	4	12	63	5	32	51	8	41	45	6	49	43	7	50	48	8	44	52	7	41	54	7	39
Peru	94	3	3	78	16	6	64	23	13	66	20	14	68	21	11	73	17	10	72	18	10	73	17	10

Table 3.2.1 Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Currently Used, by Current Age

Country	Current Age and Type of Method ^a Currently Used							Total																			
	15-19		20-24		25-29		30-34		35-39		40-44		45-49		Observed		Standardized ^b										
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E						
<i>Asia and Pacific</i>																											
Bangladesh	96	2	2	92	3	5	91	3	6	88	4	8	88	4	8	91	4	5	95	2	3	92	3	5	91	3	6
Fiji	79	4	17	68	6	26	59	6	35	50	7	43	49	7	44	55	5	40	72	2	26	59	6	35	59	6	35
Indonesia	87	2	11	73	2	25	67	4	29	66	4	30	70	4	26	77	4	19	88	2	10	74	3	23	72	4	24
Jordan	91	2	7	83	5	12	74	7	19	68	11	21	67	10	23	67	10	23	81	6	13	75	8	17	74	8	18
Korea, Rep. of	87	4	9	87	3	10	71	6	23	55	10	35	46	12	42	62	11	27	87	1	12	65	8	27	68	7	25
Malaysia	85	3	12	71	7	22	61	9	30	61	10	29	62	11	27	66	13	21	84	8	8	67	9	24	66	9	25
Nepal	100	0	0	99	0	1	98	0	2	95	0	5	95	0	5	97	0	3	98	0	2	98	0	2	97	0	3
Pakistan ^c	100	0	0	98	0	2	95	1	4	93	2	5	90	2	8	92	3	5	93	2	5	95	1	4	94	2	4
Philippines	84	11	5	71	15	14	61	21	18	57	23	20	54	25	21	64	22	14	82	11	7	64	20	16	64	20	16
Sri Lanka	86	4	10	81	7	12	70	12	18	57	16	27	59	15	26	65	16	19	80	12	8	68	13	19	69	12	19
Thailand	82	2	16	69	3	28	59	3	38	58	3	39	59	2	39	71	3	26	87	3	10	67	3	30	66	3	31
<i>Caribbean and Latin America</i>																											
Colombia	74	6	20	59	12	29	54	11	35	43	15	42	53	11	36	60	17	23	78	11	11	57	12	31	56	12	32
Costa Rica	NA	NA	NA	37	9	54	32	11	57	31	10	59	29	13	58	36	14	50	57	9	34	36	11	53	35	11	54
Dominican Rep.	87	5	8	73	5	22	59	7	34	59	6	35	60	6	34	72	8	20	82	3	15	68	6	26	67	6	27
Guyana	82	8	10	75	8	17	66	6	28	57	8	35	60	10	30	66	6	28	74	5	21	69	7	25	66	7	26
Jamaica	69	1	30	61	2	37	56	2	42	48	4	48	55	3	42	66	2	32	69	2	29	61	2	37	58	2	39
Mexico	86	3	11	73	5	22	61	8	31	62	8	30	62	11	27	75	8	17	88	4	8	70	7	23	69	7	24
Panama	NA	NA	NA	57	9	34	41	8	51	43	8	49	42	8	50	44	8	48	52	8	40	46	8	46	46	8	46
Peru	83	11	6	71	20	9	63	24	13	59	23	18	65	21	14	71	21	8	84	12	4	69	20	11	68	21	12

Table 3.2.2 Per Cent Distribution of Currently Married Women according to Type of Contraceptive Method Currently Used, by Number of Living Children

Country	Number of Living Children and Type of Method ^a Currently Used							Total																			
	None		1		2		3		4		5+		Observed		Standardized ^d												
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E									
<i>Asia and Pacific</i>																											
Bangladesh	98	1	1	94	3	3	93	3	4	91	4	5	90	4	6	88	4	8	92	3	5	92	3	5	92	3	5
Fiji	94	2	4	72	6	22	62	8	30	57	7	36	47	7	46	45	5	50	59	6	35	59	6	35	59	6	35
Indonesia	97	1	2	80	3	17	69	4	27	65	4	31	68	4	28	66	5	29	74	3	23	72	4	24	72	4	24
Jordan	96	2	2	83	5	12	73	7	20	75	8	17	74	9	17	70	9	21	75	8	17	77	7	16	77	7	16
Korea, Rep. of	92	3	5	88	3	9	61	9	30	54	11	35	54	10	36	64	8	28	65	8	27	70	7	23	70	7	23
Malaysia	94	3	3	73	6	21	65	10	25	62	11	27	60	12	28	64	10	26	67	9	24	69	9	22	69	9	22
Nepal	100	0	0	99	0	1	98	0	2	97	0	3	96	0	4	93	0	7	98	0	2	97	0	2	97	0	3
Pakistan ^c	100	0	0	99	0	1	97	1	2	95	1	4	94	2	4	90	3	7	95	1	4	94	2	4	94	2	4
Philippines	97	2	1	74	15	11	71	22	17	55	25	20	57	22	21	64	20	16	64	20	16	64	20	16	66	19	15
Sri Lanka	97	2	1	79	12	9	67	18	15	61	17	22	57	16	27	63	12	25	68	13	19	69	13	19	69	13	18
Thailand	94	1	5	69	4	27	62	2	36	58	3	39	57	3	40	69	3	28	67	3	30	67	3	30	68	3	29
<i>Caribbean and Latin America</i>																											
Colombia	87	4	9	61	11	28	50	15	35	51	10	39	49	14	37	58	13	29	57	12	31	58	12	30	58	12	30
Costa Rica ^e	80	3	17	37	11	52	27	13	60	28	12	60	32	11	57	35	11	54	36	11	53	38	10	52	38	10	52
Dominican Rep.	91	3	6	79	4	17	65	6	29	66	9	35	59	8	33	64	6	30	68	6	26	68	6	26	68	6	26
Guyana	83	8	9	79	5	16	68	8	24	66	9	25	66	9	25	59	7	34	69	7	24	68	7	24	68	7	25
Jamaica	73	2	25	69	1	30	55	2	43	60	3	37	54	4	42	52	3	45	61	2	37	59	2	37	59	2	39
Mexico	94	1	5	73	7	20	61	7	32	62	8	30	64	8	28	71	8	21	70	7	23	70	7	23	70	7	23
Panama ^e	82	5	13	54	6	40	41	10	49	40	7	53	40	7	53	45	9	46	46	8	46	49	8	44	49	8	44
Peru	93	4	3	72	20	8	61	25	14	62	22	16	65	23	12	71	19	10	69	20	11	71	19	10	71	19	10

Table 3.3.1 Per Cent Distribution of Currently Married Fecund^B Women according to Type of Contraceptive Method Currently Used, by Current Age

Country	Current Age and Type of Method ^A Currently Used												Total														
	15-19			20-24			25-29			30-34			35-39			40-44			45-49			Observed		Standardized ^B			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																											
Bangladesh	96	2	2	92	3	5	91	3	6	88	4	8	86	5	9	89	5	6	93	3	4	92	3	5	91	3	6
Fiji	79	4	17	68	6	26	59	5	36	48	8	44	44	7	49	39	7	54	41	5	54	55	6	39	55	6	39
Indonesia	87	2	11	73	2	25	66	4	30	63	5	32	62	5	33	59	7	34	65	7	28	68	4	28	67	4	29
Jordan	91	2	7	83	5	12	73	8	19	66	12	22	63	11	26	56	13	31	61	13	26	72	9	19	71	9	20
Korea, Rep. of	87	4	9	87	3	10	71	6	23	54	10	36	42	13	45	50	14	36	57	5	38	60	9	31	65	8	27
Malaysia	85	3	12	71	7	22	61	9	30	60	10	30	60	11	29	57	16	27	65	17	18	63	10	27	63	10	27
Nepal	100	0	0	99	0	1	98	0	2	95	0	5	95	0	5	96	0	4	94	0	6	97	0	3	97	0	3
Pakistan ^C	100	0	0	98	0	2	95	1	4	93	2	5	89	3	8	90	4	6	83	6	11	94	2	4	93	2	5
Philippines	84	11	5	71	15	14	61	21	18	56	23	21	52	26	22	55	28	17	60	25	15	60	22	18	61	21	17
Sri Lanka	86	4	10	80	7	13	69	12	19	55	17	28	55	17	28	54	21	25	58	26	16	63	15	22	65	14	21
Thailand	82	2	16	69	3	28	58	3	39	55	3	42	54	3	43	56	5	39	60	8	32	60	3	37	60	3	37
<i>Caribbean and Latin America</i>																											
Colombia	73	6	21	59	12	29	53	11	36	42	15	43	50	11	39	56	19	25	69	15	16	55	13	32	54	13	33
Costa Rica	NA	NA	NA	36	9	55	30	11	59	27	10	63	23	14	63	26	17	57	37	13	50	29	12	59	29	12	59
Dominican Rep.	87	5	8	72	5	23	58	7	35	58	6	36	57	7	36	66	10	24	73	5	22	66	6	28	64	6	30
Guyana	82	8	10	75	8	17	66	6	28	56	8	36	58	10	32	61	6	33	65	6	29	66	8	26	65	8	27
Jamaica	69	1	30	61	2	37	56	2	42	47	3	50	53	3	44	62	2	36	59	3	38	58	2	40	56	2	42
Mexico	86	3	11	73	5	22	60	8	32	60	8	32	58	12	30	64	11	25	69	9	22	65	8	27	65	8	27
Panama	NA	NA	NA	56	9	35	40	8	52	40	8	52	36	9	55	37	9	54	36	11	53	42	9	49	42	9	49
Peru	83	11	6	71	20	9	62	25	13	57	25	18	63	22	15	63	27	10	70	22	8	65	23	12	64	23	13

Table 3.3.2 Per Cent Distribution of Currently Married Fecund^B Women according to Type of Contraceptive Method Currently Used, by Number of Living Children

Country	Number of Living Children and Type of Method ^A Currently Used												Total											
	None			1			2			3			4			5+			Observed		Standardized ^D			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																								
Bangladesh	97	1	2	94	3	3	93	3	4	90	4	6	90	4	6	86	5	9	92	3	5	91	3	6
Fiji	93	2	5	71	6	23	59	9	32	55	7	38	44	7	49	34	6	60	55	6	39	55	6	39
Indonesia	97	1	2	77	3	20	65	4	31	59	5	36	61	5	34	55	6	39	68	4	28	65	5	30
Jordan	96	2	2	82	6	12	73	7	20	73	9	18	72	10	18	64	11	25	72	9	19	74	8	18
Korea, Rep. of	92	3	5	88	3	9	59	9	32	50	12	38	46	11	43	52	10	38	60	9	31	64	8	28
Malaysia	94	3	3	70	7	23	63	11	26	59	12	29	57	13	30	57	12	31	63	10	27	64	10	26
Nepal	100	0	0	99	0	1	98	0	2	97	0	3	96	0	4	91	0	9	97	0	3	96	0	4
Pakistan ^C	100	0	0	98	0	2	97	1	2	94	2	4	93	2	5	88	3	9	94	2	4	94	2	4
Philippines	96	3	1	73	16	11	59	23	18	53	26	21	53	24	23	56	25	19	60	22	18	62	21	17
Sri Lanka	97	2	1	78	13	9	64	19	17	56	19	25	50	19	31	52	15	33	63	15	22	63	15	22
Thailand	93	1	6	67	4	29	57	3	40	51	3	46	48	4	48	57	4	39	60	3	37	61	3	36
<i>Caribbean and Latin America</i>																								
Colombia	85	4	11	59	12	29	48	16	36	49	11	40	45	15	40	55	14	31	55	13	32	56	12	32
Costa Rica ^E	75	4	21	34	12	54	23	14	63	22	13	65	26	12	62	26	12	62	29	12	59	31	11	58
Dominican Rep.	90	3	7	78	4	18	64	6	30	54	9	37	56	9	35	60	6	34	66	6	28	65	6	29
Guyana	83	8	9	79	5	16	66	6	28	65	9	26	65	9	26	55	7	38	66	8	26	66	8	26
Jamaica	70	3	27	68	1	31	53	2	45	59	3	38	51	4	45	49	3	48	58	2	40	56	3	41
Mexico	93	1	6	71	7	22	58	8	34	59	9	32	59	9	32	65	9	26	65	8	27	66	8	26
Panama ^E	79	6	15	52	6	42	37	10	53	34	8	58	36	8	56	40	10	50	42	9	49	44	8	48
Peru	92	5	3	71	21	8	58	27	15	58	25	17	62	25	13	66	22	12	65	23	12	67	21	12

Table 3.4.1 Per Cent Distribution of Exposed^h Women according to Type of Contraceptive Method Currently Used, by Current Age

Country	Current Age and Type of Method ^a Currently Used															Total											
	15-19			20-24			25-29			30-34			35-39			40-44			45-49			Observed		Standardized ^b			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																											
Bangladesh	95	2	3	90	4	6	90	3	7	86	5	9	84	6	10	89	5	6	93	3	4	90	4	6	88	4	8
Fiji	71	5	24	58	8	34	50	7	43	43	8	49	41	8	51	38	7	55	41	5	54	48	7	45	48	7	45
Indonesia	84	3	13	67	3	30	60	5	35	57	5	38	58	6	36	57	7	36	64	7	29	63	5	32	61	5	34
Jordan	87	3	10	75	8	17	63	11	26	56	16	28	54	14	32	52	14	34	58	14	28	63	12	25	62	12	26
Korea, Rep. of	82	5	13	81	5	14	64	8	28	49	12	39	40	13	47	48	15	37	55	5	40	54	10	35	59	9	32
Malaysia	78	5	17	63	9	28	52	11	37	54	12	34	57	12	31	57	16	27	65	17	18	58	12	30	59	12	29
Nepal	100	0	0	99	0	1	97	0	3	95	0	5	94	0	6	96	0	4	94	0	6	97	0	3	96	0	4
Pakistan ^c	100	0	0	97	1	2	93	2	5	91	3	6	87	3	10	89	4	7	82	6	12	93	2	5	92	2	6
Philippines	77	16	7	60	21	19	51	27	22	48	28	24	46	29	25	52	29	19	60	25	15	52	27	21	53	26	21
Sri Lanka	82	5	13	74	10	16	64	14	22	50	19	31	52	18	30	52	22	26	58	26	16	59	17	24	60	16	24
Thailand	77	3	20	61	4	35	52	3	45	48	4	48	50	3	47	54	5	41	60	8	32	55	4	41	54	4	42
<i>Caribbean and Latin America</i>																											
Colombia	62	9	29	48	15	37	44	13	43	35	17	48	47	12	41	55	19	26	69	15	16	48	15	37	47	15	38
Costa Rica	NA	NA	NA	23	11	66	20	13	67	18	12	70	20	14	66	23	17	60	36	14	50	22	13	65	21	13	66
Dominican Rep.	80	8	12	63	7	30	49	8	43	49	7	44	53	7	40	65	10	25	73	5	22	58	8	34	58	8	34
Guyana	77	10	13	69	10	21	61	7	32	52	9	39	56	11	33	61	6	33	65	6	29	62	9	29	61	9	30
Jamaica	63	1	36	55	2	43	51	2	47	42	4	54	50	4	46	61	2	37	59	3	38	53	3	44	52	3	45
Mexico	79	4	17	64	7	29	52	10	38	53	10	37	54	13	33	61	12	27	67	10	23	58	10	32	58	10	32
Panama	NA	NA	NA	44	11	45	30	9	61	33	9	58	33	10	57	36	9	55	35	11	54	35	10	55	35	10	55
Peru	76	16	8	62	26	12	54	30	16	50	29	21	58	25	17	61	28	11	69	22	9	59	26	15	58	27	15

Table 3.4.2 Per Cent Distribution of Exposed^h Women according to Type of Contraceptive Method Currently Used, by Number of Living Children

Country	Number of Living Children and Type of Method ^a Currently Used															Total								
	None			1			2			3			4			5+			Observed		Standardized ^d			
	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E	N	I	E
<i>Asia and Pacific</i>																								
Bangladesh	97	1	2	93	3	4	92	3	5	89	5	6	89	4	7	85	5	10	90	4	6	89	4	7
Fiji	90	3	7	64	8	28	51	11	38	49	8	43	38	8	54	32	6	62	48	7	45	48	7	45
Indonesia	96	1	3	73	4	23	60	5	35	54	6	40	57	6	37	49	7	44	63	5	32	60	5	35
Jordan	93	4	3	74	8	18	63	10	27	64	12	24	63	13	24	57	13	30	63	12	25	65	11	24
Korea, Rep. of	84	7	9	83	4	13	53	11	36	46	13	41	43	12	45	51	10	39	54	10	35	59	9	32
Malaysia	90	5	5	63	8	29	57	13	30	55	13	32	53	14	33	55	12	33	58	12	30	59	12	29
Nepal	100	0	0	99	0	1	98	0	2	97	0	3	95	0	5	91	0	9	97	0	3	96	0	4
Pakistan ^c	100	0	0	98	0	2	95	2	3	93	2	5	91	3	6	86	4	10	93	2	5	92	2	6
Philippines	95	4	1	64	21	15	51	27	22	45	30	25	46	28	26	50	28	22	52	27	21	55	25	20
Sri Lanka	96	2	2	74	15	11	59	22	19	52	21	27	47	20	33	49	16	35	59	17	24	58	17	25
Thailand	90	1	9	62	5	33	53	3	44	47	4	49	42	4	54	53	4	43	55	4	41	55	4	41
<i>Caribbean and Latin America</i>																								
Colombia	76	7	17	50	14	36	40	18	42	43	12	45	41	16	43	51	15	34	48	15	37	49	14	37
Costa Rica ^e	61	6	33	25	13	62	18	14	68	17	14	69	20	13	67	22	13	65	22	13	65	24	13	64
Dominican Rep.	85	4	11	70	6	24	56	7	37	49	10	41	50	10	40	55	7	38	58	8	34	58	8	35
Guyana	78	10	12	74	6	20	63	9	28	61	11	28	62	10	28	53	8	39	62	9	29	62	9	29
Jamaica	66	3	31	64	1	35	48	2	50	54	3	43	46	5	49	46	3	51	53	3	44	52	3	45
Mexico	87	3	10	62	9	29	51	9	40	53	10	37	54	11	35	59	11	30	58	10	32	59	9	31
Panama ^e	69	9	22	42	8	50	30	11	59	27	9	64	29	9	62	37	10	53	35	10	55	37	10	53
Peru	84	9	7	64	26	10	51	31	18	52	28	20	57	29	14	63	24	13	59	26	15	61	25	14

GROUP 4 CURRENT USE BY INDIVIDUAL METHODS

LIST OF TABLES

4.1.1.1 Percentage of Currently Married Women Currently Using the Pill, by Current Age	52	4.1.1.2 Percentage of Currently Married Women Currently Using the Pill, by Number of Living Children	58
4.1.2.1 Percentage of Currently Married Women Currently Using IUD, by Current Age	52	4.1.2.2 Percentage of Currently Married Women Currently Using IUD, by Number of Living Children	58
4.1.3.1 Percentage of Currently Married Women Currently Using Other Female Scientific Methods, by Current Age	53	4.1.3.2 Percentage of Currently Married Women Currently Using Other Female Scientific Methods, by Number of Living Children	59
4.1.4.1 Percentage of Currently Married Women Currently Using Douche, by Current Age	53	4.1.4.2 Percentage of Currently Married Women Currently Using Douche, by Number of Living Children	59
4.1.5.1 Percentage of Currently Married Women Currently Using Condom, by Current Age	54	4.1.5.2 Percentage of Currently Married Women Currently Using Condom, by Number of Living Children	60
4.1.6.1 Percentage of Currently Married Women Currently Using Rhythm, by Current Age	54	4.1.6.2 Percentage of Currently Married Women Currently Using Rhythm, by Number of Living Children	60
4.1.7.1 Percentage of Currently Married Women Currently Using Withdrawal, by Current Age	55	4.1.7.2 Percentage of Currently Married Women Currently Using Withdrawal by Number of Living Children	61
4.1.8.1 Percentage of Currently Married Women Currently Using Abstinence, by Current Age	55	4.1.8.2 Percentage of Currently Married Women Currently Using Abstinence, by Number of Living Children	61
4.1.9.1 Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Current Age	56	4.1.9.2 Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Number of Living Children	62
4.1.10.1 Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Current Age	56	4.1.10.2 Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Number of Living Children	62
4.1.11.1 Percentage of Currently Married Women Currently Using Injection, by Current Age	57	4.1.11.2 Percentage of Currently Married Women Currently Using Injection, by Number of Living Children	63
4.1.12.1 Percentage of Currently Married Women Currently Using Other Methods, by Current Age	57	4.1.12.2 Percentage of Currently Married Women Currently Using Other Methods, by Number of Living Children	63

4.2.1.1 Percentage of Exposed Women Currently Using the Pill, by Current Age	64	4.2.1.2 Percentage of Exposed Women Currently Using the Pill, by Number of Living Children	70
4.2.2.1 Percentage of Exposed Women Currently Using IUD, by Current Age	64	4.2.2.2 Percentage of Exposed Women Currently Using IUD, by Number of Living Children	70
4.2.3.1 Percentage of Exposed Women Currently Using Other Female Scientific Methods, by Current Age	65	4.2.3.2 Percentage of Exposed Women Currently Using Other Female Scientific Methods, by Number of Living Children	71
4.2.4.1 Percentage of Exposed Women Currently Using Douche, by Current Age	65	4.2.4.2 Percentage of Exposed Women Currently Using Douche, by Number of Living Children	71
4.2.5.1 Percentage of Exposed Women Currently Using Condom, by Current Age	66	4.2.5.2 Percentage of Exposed Women Currently Using the Condom, by Number of Living Children	72
4.2.6.1 Percentage of Exposed Women Currently Using Rhythm, by Current Age	66	4.2.6.2 Percentage of Exposed Women Currently Using Rhythm, by Number of Living Children	72
4.2.7.1 Percentage of Exposed Women Currently Using Withdrawal, by Current Age	67	4.2.7.2 Percentage of Exposed Women Currently Using Withdrawal, by Number of Living Children	73
4.2.8.1 Percentage of Exposed Women Currently Using Abstinence, by Current Age	67	4.2.8.2 Percentage of Exposed Women Currently Using Abstinence, by Number of Living Children	73
4.2.9.1 Percentage of Exposed Women Sterilized for Contraceptive Purposes, by Current Age	68	4.2.9.2 Percentage of Exposed Women Sterilized for Contraceptive Purposes, by Number of Living Children	74
4.2.10.1 Percentage of Exposed Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Current Age	68	4.2.10.2 Percentage of Exposed Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Number of Living Children	74
4.2.11.1 Percentage of Exposed Women Currently Using Injection, by Current Age	69	4.2.11.2 Percentage of Exposed Women Currently Using Injection, by Number of Living Children	75
4.2.12.1 Percentage of Exposed Women Currently Using Other Methods, by Current Age	69	4.2.12.2 Percentage of Exposed Women Currently Using Other Methods, by Number of Living Children	75

Table 4.1.1.1 Percentage of Currently Married Women Currently Using the Pill, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.1	3.0	4.6	5.2	2.5	1.8	1.0	2.9
Fiji	6.7	11.4	10.9	7.8	7.8	5.0	1.5	8.2
Indonesia	8.8	18.4	19.7	18.8	14.6	10.3	5.3	14.9
Jordan	5.7	10.0	15.0	13.0	14.1	14.1	6.3	11.9
Korea, Rep. of	7.3	5.2	7.1	10.8	14.0	6.9	1.7	8.4
Malaysia	11.6	19.1	24.0	18.2	15.9	10.7	3.0	16.2
Nepal	0.0	0.2	0.3	1.1	0.5	0.4	0.0	0.4
Pakistan ^c	0.0	0.2	1.1	1.9	2.0	1.1	0.2	1.0
Philippines	2.6	6.0	6.1	5.6	4.7	2.7	1.1	4.5
Sri Lanka	1.1	2.1	2.1	2.1	1.9	0.4	0.2	1.5
Thailand	13.3	15.5	21.2	15.7	13.4	8.5	2.1	13.7
<i>Caribbean and Latin America</i>								
Colombia	12.2	17.3	18.1	16.6	9.6	6.8	4.7	13.3
Costa Rica	NA	38.4	31.2	24.7	14.3	10.0	2.9	22.5
Dominican Rep.	4.7	12.8	12.4	6.8	5.0	2.3	2.5	7.9
Guyana	8.9	10.0	14.7	11.7	6.3	3.6	0.7	9.0
Jamaica	12.2	15.2	19.9	12.5	8.3	5.4	2.5	11.9
Mexico	7.8	13.5	14.9	13.1	10.2	5.9	2.7	10.8
Panama	NA	26.1	27.3	16.2	10.8	6.9	1.8	17.0
Peru	3.2	4.2	7.5	5.9	3.4	1.7	0.6	4.2

Table 4.1.2.1 Percentage of Currently Married Women Currently Using IUD, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.2	0.0	0.6	1.7	0.5	0.5	0.4
Fiji	0.4	4.2	5.4	6.1	5.0	5.0	2.3	4.7
Indonesia	1.1	4.6	5.8	8.0	8.0	6.2	3.4	5.6
Jordan	0.4	0.6	2.5	3.6	2.6	2.4	0.9	2.0
Korea, Rep. of	1.8	2.5	6.7	10.0	12.5	9.1	3.4	8.0
Malaysia	0.0	0.2	0.6	1.4	0.8	1.2	0.2	0.7
Nepal	0.0	0.1	0.0	0.1	0.1	0.0	0.2	0.1
Pakistan ^c	0.0	0.2	0.7	0.7	1.6	0.7	0.8	0.6
Philippines	0.4	1.8	2.4	3.0	3.0	2.3	1.3	2.3
Sri Lanka	5.4	5.6	4.7	6.4	5.0	4.9	0.9	4.7
Thailand	1.0	6.8	5.2	10.7	7.4	4.4	1.3	5.9
<i>Caribbean and Latin America</i>								
Colombia	3.9	6.3	11.2	12.5	11.0	6.2	2.5	8.5
Costa Rica	NA	4.8	5.1	6.3	6.7	4.3	3.3	5.2
Dominican Rep.	0.0	2.0	5.7	3.8	3.0	1.7	0.6	2.8
Guyana	0.9	5.9	8.3	9.1	5.3	2.2	2.6	5.6
Jamaica	0.8	0.8	1.5	3.4	3.8	1.8	2.5	2.0
Mexico	1.6	4.7	9.2	7.4	7.0	3.6	0.6	5.7
Panama	NA	3.2	5.1	5.0	3.0	2.4	1.5	3.7
Peru	0.4	1.4	1.4	2.9	1.0	1.1	0.2	1.3

Table 4.1.3.1 Percentage of Currently Married Women Currently Using Other Female Scientific Methods, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.0	0.1	0.0	0.0	0.2	0.2	0.0
Fiji	0.0	0.0	0.1	0.0	0.0	0.2	0.0	0.0
Indonesia	0.0	0.0	0.1	0.2	0.1	0.0	0.0	0.1
Jordan	0.0	0.2	0.1	0.5	0.0	0.0	0.0	0.2
Korea, Rep. of	0.0	0.0	0.4	0.6	0.1	0.4	0.0	0.3
Malaysia	0.0	0.0	0.1	0.2	0.1	0.3	0.3	0.1
Nepal	N Ask							
Pakistan ^c	0.0	0.2	0.1	0.1	0.4	0.2	0.2	0.2
Philippines	0.0	0.1	0.0	0.1	0.3	0.2	0.2	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Thailand	0.0	0.2	0.0	0.2	0.2	0.0	0.0	0.1
<i>Caribbean and Latin America</i>								
Colombia	1.7	2.1	1.7	3.8	3.0	2.5	0.7	2.3
Costa Rica	NA	1.2	1.8	2.2	2.0	2.0	1.0	1.7
Dominican Rep.	1.2	1.5	1.7	2.3	2.3	0.6	0.6	1.6
Guyana	0.0	0.3	0.5	0.6	0.0	0.3	0.3	0.3
Jamaica	1.6	1.7	0.5	3.4	1.3	0.7	1.3	1.5
Mexico	0.5	1.2	1.7	2.5	1.2	0.6	0.9	1.4
Panama	NA	1.2	2.8	1.9	2.4	0.9	0.4	1.8
Peru	0.4	0.5	0.6	1.7	1.3	0.8	0.0	0.8

Table 4.1.4.1 Percentage of Currently Married Women Currently Using Douche, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.0	0.2	0.3	0.0	0.0	0.0	0.1
Fiji	N Ask							
Indonesia	0.4	0.0	0.0	0.1	0.0	0.0	0.1	0.1
Jordan	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0
Korea, Rep. of	0.0	0.0	0.0	0.5	0.5	0.4	0.0	0.3
Malaysia	N Ask							
Nepal								
Pakistan								
Philippines	0.5	0.1	0.1	0.2	0.1	0.2	0.2	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Thailand	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.1
<i>Caribbean and Latin America</i>								
Colombia	1.1	0.0	0.5	1.1	0.2	0.8	0.4	0.5
Costa Rica	NA	0.6	0.4	0.0	0.2	0.3	0.3	0.3
Dominican Rep.	1.8	0.0	0.6	1.1	0.7	0.0	0.0	0.6
Guyana	2.5	1.2	1.6	2.8	2.3	2.8	0.3	1.9
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.2	0.1	0.0	0.2	0.7	0.2	0.0	0.2
Panama	NA	1.0	0.7	0.3	0.7	0.9	0.7	0.7
Peru	2.6	3.4	2.9	4.0	4.0	3.2	2.6	3.3

Table 4.1.5.1 Percentage of Currently Married Women Currently Using Condom, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.9	0.9	0.6	0.8	0.7	0.9	0.5	0.7
Fiji	9.4	8.9	8.8	5.8	3.3	2.5	1.0	6.0
Indonesia	0.7	1.4	2.6	2.2	2.2	2.5	0.4	1.8
Jordan	0.6	0.7	0.8	1.7	2.7	1.9	1.2	1.4
Korea, Rep. of	0.0	2.3	6.4	7.1	7.0	3.6	0.8	5.1
Malaysia	0.4	1.7	3.6	3.8	3.6	2.3	0.8	2.7
Nepal	0.4	0.4	0.3	0.1	0.2	0.0	0.0	0.2
Pakistan ^c	0.0	1.0	1.7	1.3	1.6	0.6	0.3	1.0
Philippines	1.8	4.1	4.1	4.2	4.4	3.3	0.8	3.6
Sri Lanka	3.2	2.2	3.2	3.4	3.0	0.6	0.1	2.3
Thailand	0.0	0.7	0.3	0.5	0.5	0.6	0.0	0.4
<i>Caribbean and Latin America</i>								
Colombia	1.7	1.9	1.4	1.7	3.7	0.6	0.4	1.7
Costa Rica	NA	8.6	9.5	9.6	9.2	8.3	6.8	8.8
Dominican Rep.	1.2	1.5	0.6	2.3	1.7	2.9	1.2	1.5
Guyana	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0
Jamaica	10.6	10.4	7.2	5.2	5.4	2.5	1.7	6.6
Mexico	0.2	0.2	1.3	0.7	1.0	0.8	0.8	0.8
Panama	NA	1.6	1.3	1.2	1.4	1.2	0.4	1.2
Peru	0.0	1.0	0.6	1.9	1.3	1.4	0.4	1.1

Table 4.1.6.1 Percentage of Currently Married Women Currently Using Rhythm, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.3	1.2	1.3	1.7	0.9	2.0	0.5	1.1
Fiji	0.9	1.9	2.0	3.9	2.2	2.5	1.3	2.3
Indonesia	0.0	0.4	1.4	1.2	1.2	0.6	0.5	0.8
Jordan	0.3	1.5	1.9	4.0	2.5	2.5	0.9	2.1
Korea, Rep. of	3.6	1.3	3.9	6.6	5.6	6.3	1.3	4.6
Malaysia	0.4	2.7	3.8	4.9	3.8	4.0	1.5	3.5
Nepal	N Ask							
Pakistan ^c	0.0	0.0	0.2	0.2	0.3	0.1	0.0	0.1
Philippines	3.5	5.5	9.1	10.6	11.3	9.4	4.0	8.5
Sri Lanka	2.8	3.8	7.9	11.2	10.2	9.0	5.1	8.0
Thailand	0.9	1.2	0.7	0.7	0.7	1.1	0.7	0.9
<i>Caribbean and Latin America</i>								
Colombia	2.8	6.1	4.1	4.7	5.4	6.8	5.1	5.1
Costa Rica	NA	3.4	4.2	5.7	6.5	7.1	4.2	5.1
Dominican Rep.	0.0	1.3	1.7	1.5	1.0	1.7	0.6	1.2
Guyana	1.5	3.8	2.7	2.8	5.3	1.7	0.7	2.9
Jamaica	0.0	0.2	0.2	0.6	0.3	0.7	0.0	0.3
Mexico	0.5	2.2	4.3	3.0	5.0	3.3	0.9	3.1
Panama	NA	3.6	2.1	2.2	2.3	2.4	2.6	2.5
Peru	4.4	12.4	14.4	13.3	9.9	10.1	4.8	10.9

Table 4.1.7.1 Percentage of Currently Married Women Currently Using Withdrawal, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.5	0.7	0.4	0.2	0.8	0.2	0.7	0.5
Fiji	1.8	3.9	2.8	2.7	3.6	1.6	1.0	2.8
Indonesia	0.5	0.4	0.3	0.3	0.3	0.4	0.0	0.3
Jordan	1.6	1.8	3.9	4.1	3.5	4.5	3.2	3.3
Korea, Rep. of	0.0	1.6	2.0	3.0	4.3	3.3	0.0	2.6
Malaysia	1.7	1.1	1.7	1.9	2.1	3.3	1.5	1.9
Nepal	N Ask							
Pakistan ^c	0.0	0.1	0.0	0.3	0.1	0.0	0.0	0.1
Philippines	6.6	8.7	9.9	10.2	11.0	10.2	5.8	9.4
Sri Lanka	0.5	1.7	2.1	2.1	1.1	1.0	0.5	1.5
Thailand	1.0	0.7	1.3	0.7	0.6	1.2	0.9	0.9
<i>Caribbean and Latin America</i>								
Colombia	1.7	3.6	3.9	6.8	3.5	8.5	4.4	4.7
Costa Rica	NA	4.6	5.3	3.9	5.4	5.1	2.9	4.6
Dominican Rep.	2.9	3.6	4.0	3.0	4.7	5.2	1.9	3.7
Guyana	0.9	1.5	0.8	0.8	0.9	0.6	1.3	1.0
Jamaica	0.8	1.5	1.7	2.1	1.3	1.1	1.3	1.4
Mexico	1.8	2.8	3.4	4.9	4.8	4.2	2.3	3.6
Panama	NA	2.0	3.0	2.9	3.7	3.0	3.7	3.0
Peru	2.4	2.6	3.9	4.1	3.9	3.2	1.7	3.3

Table 4.1.8.1 Percentage of Currently Married Women Currently Using Abstinence, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.0	1.2	0.4	1.6	2.4	1.5	1.1	1.2
Fiji	0.0	0.2	0.2	0.2	0.1	0.9	0.0	0.3
Indonesia	0.4	0.8	0.7	1.0	1.3	1.6	1.3	1.0
Jordan	0.0	0.4	0.4	0.3	0.7	0.0	0.9	0.4
Korea, Rep. of	0.0	0.2	0.0	0.1	1.2	0.5	0.2	0.4
Malaysia	0.0	0.7	0.6	1.4	2.0	3.9	3.4	1.7
Nepal	0.0	0.1	0.0	0.0	0.3	0.0	0.0	0.0
Pakistan ^c	0.1	0.2	0.8	1.4	1.7	2.7	1.9	1.2
Philippines	0.4	1.3	2.1	2.1	2.2	2.2	1.1	1.8
Sri Lanka	0.3	1.6	1.7	3.0	3.6	5.8	6.8	3.5
Thailand	0.5	0.5	0.4	0.7	1.0	0.8	0.9	0.7
<i>Caribbean and Latin America</i>								
Colombia	0.0	0.9	1.0	1.3	0.7	1.1	0.7	0.9
Costa Rica	NA	0.2	0.5	0.0	0.4	0.9	1.6	0.5
Dominican Rep.	N Ask							
Guyana	1.5	1.4	0.9	1.2	0.9	0.6	1.6	1.2
Jamaica	0.0	0.4	0.0	0.3	1.3	0.0	0.8	0.4
Mexico	N Ask							
Panama	NA	1.8	1.3	1.7	1.2	1.2	1.1	1.4
Peru	0.8	1.1	2.2	1.5	2.0	4.2	2.3	2.1

Table 4.1.9.1 Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.1	0.3	0.6	0.9	0.6	0.2	0.3
Fiji	0.0	0.9	9.9	22.1	27.8	26.6	20.7	15.8
Indonesia	0.0	0.0	0.1	0.3	0.7	0.4	0.3	0.3
Jordan	0.0	0.0	0.4	1.6	3.6	4.6	3.5	1.8
Korea, Rep. of	0.0	0.0	0.3	3.3	2.5	2.0	1.9	1.7
Malaysia	0.0	0.3	1.2	4.9	6.1	5.7	3.5	3.4
Nepal	0.0	0.1	0.0	0.0	0.4	0.0	0.0	0.1
Pakistan ^c	0.0	0.0	0.4	0.7	2.0	1.7	3.0	0.9
Philippines	0.0	1.4	4.5	6.3	7.2	5.3	2.7	4.7
Sri Lanka	0.5	1.7	7.7	13.1	14.5	11.6	6.2	9.2
Thailand	0.5	1.5	6.5	8.6	9.9	8.6	4.7	6.3
<i>Caribbean and Latin America</i>								
Colombia	0.0	0.8	2.2	6.6	8.2	6.2	2.9	4.0
Costa Rica	NA	0.6	6.8	13.1	23.0	20.3	15.3	12.3
Dominican Rep.	0.0	3.8	13.5	19.9	21.7	12.7	8.7	11.9
Guyana	0.0	0.3	3.0	12.9	17.0	19.8	14.5	8.5
Jamaica	0.0	0.6	2.5	13.4	16.2	19.1	10.1	8.1
Mexico	0.2	0.5	1.3	3.6	5.0	5.3	3.4	2.7
Panama	NA	0.8	13.6	23.8	31.4	35.8	35.5	21.2
Peru	0.0	0.2	1.3	4.2	6.3	2.9	2.9	2.8

Table 4.1.10.1 Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.3	0.3	0.4	2.3	0.7	0.4	0.5
Fiji	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.1
Indonesia	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Jordan	0.0	0.0	0.0	0.0	0.1	0.1	0.9	0.1
Korea, Rep. of	0.0	0.2	1.4	2.8	6.0	5.5	4.0	3.3
Malaysia	0.0	0.0	0.2	0.2	0.7	0.8	0.7	0.4
Nepal	0.0	0.3	1.6	3.2	3.3	2.1	1.7	1.6
Pakistan ^c	0.0	0.0	0.0	0.1	0.2	0.2	0.0	0.1
Philippines	0.0	0.3	0.6	0.9	1.1	0.5	0.4	0.7
Sri Lanka	0.0	0.2	0.7	1.3	1.0	0.9	0.0	0.7
Thailand	0.0	0.9	1.9	1.6	4.9	2.6	1.4	2.1
<i>Caribbean and Latin America</i>								
Colombia	0.0	0.2	0.3	0.2	0.0	0.3	0.0	0.2
Costa Rica	NA	0.0	1.4	2.0	0.7	0.6	1.0	1.0
Dominican Rep.	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.1
Guyana	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.0	0.0	0.1	0.3	0.7	0.2	0.0	0.2
Panama	NA	0.2	0.0	0.9	0.5	0.3	0.3	0.4
Peru	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 4.1.11.1 Percentage of Currently Married Women Currently Using Injection, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	N Ask							
Fiji	0.9	0.7	0.1	0.3	0.0	0.2	0.0	0.3
Indonesia	0.2	0.2	0.5	0.3	0.2	0.0	0.2	0.2
Jordan	N Ask							
Korea, Rep. of	0.0	0.0	0.3	0.3	0.3	0.1	0.0	0.2
Malaysia	0.0	0.3	0.3	0.4	0.2	0.1	0.0	0.2
Nepal	N Ask							
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Philippines	0.3	0.2	0.3	0.2	0.2	0.0	0.0	0.2
Sri Lanka	0.0	0.7	0.1	0.6	0.3	0.1	0.2	0.3
Thailand	0.9	2.1	2.8	1.5	2.6	1.6	0.5	1.9
<i>Caribbean and Latin America</i>								
Colombia	1.1	0.4	0.2	0.4	0.9	0.3	0.0	0.4
Costa Rica	NA	0.6	1.6	1.2	2.0	4.0	3.9	2.0
Dominican Rep.	0.6	0.3	0.0	0.0	0.0	0.0	0.6	0.2
Guyana	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Jamaica	4.5	8.4	9.5	9.5	5.4	1.1	0.4	6.2
México	0.9	1.6	2.1	2.4	2.2	1.2	0.0	1.7
Panama	NA	1.4	0.8	0.2	0.5	0.3	0.4	0.6
Peru	1.7	1.8	1.5	0.9	0.8	0.2	0.4	1.0

Table 4.1.12.1 Percentage of Currently Married Women Currently Using Other Methods, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.0	0.3	0.4	0.3	0.5	0.0	0.2
Fiji	0.9	0.2	0.5	0.3	0.7	0.4	0.0	0.4
Indonesia	0.9	0.7	1.7	1.5	1.6	1.3	0.7	1.2
Jordan	0.3	1.7	1.3	2.9	3.3	2.9	1.3	2.0
Korea, Rep. of	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Malaysia	1.2	2.9	2.7	1.9	2.5	1.7	1.3	2.2
Nepal	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pakistan ^c	0.0	0.0	0.2	0.0	0.3	0.0	0.3	0.1
Philippines	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.1
Sri Lanka	0.0	0.0	0.0	0.1	0.0	0.2	0.0	0.1
Thailand	0.0	0.6	0.3	0.6	0.0	0.0	0.0	0.3
<i>Caribbean and Latin America</i>								
Colombia	0.6	1.1	1.4	0.8	0.9	0.0	0.0	0.8
Costa Rica	NA	0.4	0.5	0.2	0.2	0.9	0.0	0.4
Dominican Rep.	0.6	0.0	0.3	0.0	0.0	1.2	0.6	0.3
Guyana	1.2	0.0	0.3	0.6	0.2	0.0	0.7	0.4
Jamaica	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0
Mexico	0.5	0.1	0.3	0.0	0.2	0.0	0.2	0.2
Panama	NA	0.2	0.7	0.3	0.5	0.3	0.0	0.4
Peru	0.7	0.7	0.8	0.6	0.6	0.6	0.2	0.6

Table 4.1.1.2 Percentage of Currently Married Women Currently Using the Pill, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.3	1.7	3.2	3.3	3.3	4.5	2.9
Fiji	1.6	11.0	11.9	12.9	6.7	6.3	8.2
Indonesia	1.9	14.4	19.9	19.6	15.9	15.9	14.9
Jordan	1.3	8.7	14.7	12.9	12.7	13.2	11.9
Korea, Rep. of	3.3	3.6	8.5	10.6	11.0	8.8	8.4
Malaysia	1.3	17.0	18.8	19.4	19.7	15.7	16.2
Nepal	0.0	0.3	0.0	0.5	0.5	1.2	0.4
Pakistan ^c	0.0	0.3	0.2	0.9	1.3	2.0	1.0
Philippines	0.2	5.0	6.6	5.8	4.9	3.6	4.5
Sri Lanka	0.3	1.7	2.2	2.0	2.0	1.2	1.5
Thailand	3.7	19.2	19.1	15.0	11.8	10.7	13.7
<i>Caribbean and Latin America</i>							
Colombia	5.2	17.0	15.9	16.7	14.0	10.9	13.3
Costa Rica ^e	8.8	35.7	32.8	21.5	20.7	14.0	22.5
Dominican Rep.	3.6	11.7	12.7	10.2	6.4	5.2	7.9
Guyana	7.0	12.4	14.1	10.5	9.6	5.5	9.0
Jamaica	8.9	14.9	17.8	16.1	9.6	6.9	11.9
Mexico	2.7	11.9	16.3	13.4	13.9	8.4	10.8
Panama ^e	9.2	27.7	27.6	18.7	15.9	8.8	17.0
Peru	1.7	4.4	6.8	5.4	4.0	2.7	4.2

Table 4.1.2.2 Percentage of Currently Married Women Currently Using IUD, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.1	0.0	0.5	0.2	0.9	0.4
Fiji	0.0	1.7	4.8	4.8	9.7	5.8	4.7
Indonesia	0.0	2.3	5.5	7.5	8.4	9.1	5.6
Jordan	0.0	0.9	2.6	1.0	2.2	2.6	2.0
Korea, Rep. of	0.0	1.4	8.7	9.9	11.4	9.5	8.0
Malaysia	0.2	0.5	0.9	1.0	0.6	0.7	0.7
Nepal	0.0	0.0	0.1	0.1	0.0	0.2	0.1
Pakistan ^c	0.1	0.2	0.3	0.6	0.5	1.4	0.6
Philippines	0.1	1.2	2.7	3.0	3.6	2.2	2.3
Sri Lanka	0.0	3.4	5.9	6.2	6.6	4.7	4.7
Thailand	0.0	3.2	7.0	9.6	7.6	5.8	5.9
<i>Caribbean and Latin America</i>							
Colombia	0.0	6.6	10.9	12.1	13.3	7.3	8.5
Costa Rica ^e	1.1	4.0	5.2	6.7	6.0	5.7	5.2
Dominican Rep.	0.0	0.8	3.9	4.9	3.4	3.3	2.8
Guyana	0.0	1.7	6.6	7.3	7.1	7.9	5.6
Jamaica	1.0	0.2	1.6	1.7	3.7	3.4	2.0
Mexico	0.2	4.7	7.3	7.9	7.4	5.2	5.7
Panama ^e	0.0	3.6	5.3	3.7	4.9	3.2	3.7
Peru	0.0	1.5	2.8	0.9	1.4	1.0	1.3

Table 4.1.3.2 Percentage of Currently Married Women Currently Using Other Female Scientific Methods, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Fiji	0.0	0.2	0.0	0.2	0.0	0.0	0.0
Indonesia	0.0	0.1	0.0	0.0	0.1	0.1	0.1
Jordan	0.0	0.4	0.3	0.3	0.0	0.1	0.2
Korea, Rep.of	0.0	0.4	0.1	0.5	0.3	0.2	0.3
Malaysia	0.0	0.0	0.1	0.3	0.3	0.1	0.1
Nepal	N Ask						
Pakistan ^c	0.0	0.2	0.0	0.3	0.0	0.3	0.2
Philippines	0.0	0.0	0.1	0.2	0.2	0.2	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Thailand	0.0	0.0	0.2	0.0	0.4	0.0	0.1
<i>Caribbean and Latin America</i>							
Colombia	0.8	2.0	4.1	1.9	2.9	2.0	2.3
Costa Rica ^e	1.1	1.0	2.9	3.5	1.1	0.9	1.7
Dominican Rep.	1.3	1.1	2.3	1.2	0.5	2.1	1.6
Guyana	0.0	0.0	0.2	0.2	0.8	0.5	0.3
Jamaica	2.3	1.2	1.6	2.0	0.0	1.5	1.5
Mexico	1.0	0.7	2.4	2.2	2.1	0.7	1.4
Panama ^e	1.2	2.7	2.3	3.2	0.6	1.0	1.8
Peru	0.8	0.3	1.2	1.1	0.9	0.8	0.8

Table 4.1.4.2 Percentage of Currently Married Women Currently Using Douche, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.0	0.2	0.0	0.3	0.1	0.1
Fiji	N Ask						
Indonesia	0.0	0.2	0.1	0.0	0.0	0.0	0.1
Jordan	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Korea, Rep.of	0.0	0.0	0.3	0.4	0.3	0.2	0.3
Malaysia	N Ask						
Nepal							
Pakistan							
Philippines	0.0	0.4	0.2	0.0	0.3	0.1	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Thailand	0.0	0.0	0.1	0.0	0.2	0.0	0.1
<i>Caribbean and Latin America</i>							
Colombia	0.4	0.0	1.0	1.2	0.0	0.4	0.5
Costa Rica ^e	0.0	0.2	0.4	0.5	0.4	0.2	0.3
Dominican Rep.	0.9	0.0	0.8	0.4	1.5	0.3	0.6
Guyana	1.6	0.6	1.4	2.7	3.1	2.1	1.9
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.0	0.1	0.4	0.3	0.1	0.2	0.2
Panama ^e	1.2	0.6	0.6	0.6	1.1	0.5	0.7
Peru	1.5	2.6	3.5	3.1	3.5	3.9	3.3

Table 4.1.5.2 Percentage of Currently Married Women Currently Using Condom, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.7	0.8	0.4	0.9	0.8	0.8	0.7
Fiji	2.0	7.8	8.4	9.7	6.5	3.9	6.0
Indonesia	0.0	0.8	1.5	3.0	2.5	3.0	1.8
Jordan	0.0	1.7	1.6	2.1	1.2	1.4	1.4
Korea, Rep. of	1.1	2.8	7.7	7.0	6.0	3.7	5.1
Malaysia	1.3	3.2	3.4	3.7	2.6	2.2	2.7
Nepal	0.1	0.1	0.6	0.0	0.1	0.4	0.2
Pakistan ^c	0.0	0.6	0.9	1.3	1.2	1.6	1.0
Philippines	0.1	3.6	4.4	3.6	4.1	3.5	3.6
Sri Lanka	1.0	3.2	3.6	3.4	2.4	1.0	2.3
Thailand	0.3	0.3	0.9	0.2	0.9	0.2	0.4
<i>Caribbean and Latin America</i>							
Colombia	1.6	2.2	1.9	2.2	1.3	1.3	1.7
Costa Rica ^e	3.8	9.0	13.5	11.1	9.1	5.8	8.8
Dominican Rep.	0.4	1.9	1.9	1.2	2.5	1.5	1.5
Guyana	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Jamaica	9.2	7.2	9.8	5.4	7.3	3.4	6.6
Mexico	0.0	0.3	0.7	1.4	0.9	0.8	0.8
Panama ^e	1.2	2.1	1.5	1.1	1.9	0.7	1.2
Peru	0.0	0.5	1.2	2.0	1.1	0.9	1.1

Table 4.1.6.2 Percentage of Currently Married Women Currently Using Rhythm, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.3	0.8	0.5	2.1	1.2	1.3	1.1
Fiji	0.9	0.9	3.1	3.1	3.6	2.3	2.3
Indonesia	0.2	0.4	1.1	0.6	1.3	1.2	0.8
Jordan	0.3	1.3	4.7	2.1	4.4	1.5	2.1
Korea, Rep. of	3.1	1.7	5.5	6.6	5.8	3.5	4.6
Malaysia	1.1	2.7	5.1	5.6	4.4	2.4	3.5
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.0	0.1	0.7	0.0	0.1
Philippines	1.3	6.7	9.7	11.3	10.4	8.0	8.5
Sri Lanka	1.1	8.2	12.4	10.1	12.2	5.1	8.0
Thailand	0.3	1.5	1.2	0.6	0.9	0.6	0.9
<i>Caribbean and Latin America</i>							
Colombia	2.0	7.4	6.8	3.4	7.8	3.9	5.1
Costa Rica ^e	2.2	4.3	6.9	6.4	7.0	3.8	5.1
Dominican Rep.	1.3	0.8	0.8	2.8	0.5	1.1	1.2
Guyana	3.5	1.7	3.6	3.2	3.4	2.5	2.9
Jamaica	0.3	0.0	0.0	1.3	0.9	0.0	0.3
Mexico	0.8	2.6	4.4	4.5	3.8	2.5	3.1
Panama ^e	1.7	3.0	4.1	2.8	1.9	1.9	2.5
Peru	1.7	13.9	15.4	13.7	10.9	7.7	10.9

Table 4.1.7.2 Percentage of Currently Married Women Currently Using Withdrawal, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.7	0.7	0.6	0.6	0.4	0.5
Fiji	0.7	4.4	4.1	3.4	3.1	1.9	2.8
Indonesia	0.0	0.4	0.4	0.2	0.4	0.4	0.3
Jordan	1.3	2.2	1.4	4.5	2.5	4.2	3.3
Korea, Rep. of	0.0	1.3	2.7	3.8	2.6	2.9	2.6
Malaysia	1.1	1.1	1.5	2.0	3.0	2.2	1.9
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.3	0.0	0.1	0.1	0.1
Philippines	0.9	7.0	10.3	11.0	10.2	10.1	9.4
Sri Lanka	0.4	2.0	1.9	2.3	0.9	1.1	1.5
Thailand	0.3	1.3	0.5	0.7	1.3	1.0	0.9
<i>Caribbean and Latin America</i>							
Colombia	0.4	2.7	4.3	4.6	4.5	7.1	4.7
Costa Rica ^e	1.1	5.5	5.2	4.7	3.5	4.9	4.6
Dominican Rep.	0.4	3.0	3.9	5.3	5.4	3.9	3.7
Guyana	1.2	1.3	1.1	1.2	0.8	0.7	1.0
Jamaica	1.6	1.0	1.3	0.3	2.7	1.8	1.4
Mexico	0.0	3.9	2.3	3.0	4.0	5.0	3.6
Panama ^e	1.7	1.5	2.4	2.6	2.7	4.4	3.0
Peru	0.0	2.3	4.0	3.9	4.6	3.1	3.3

Table 4.1.8.2 Percentage of Currently Married Women Currently Using Abstinence, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.9	1.3	0.9	1.0	2.0	1.2
Fiji	0.0	0.2	0.3	0.2	0.0	0.5	0.3
Indonesia	0.1	0.8	0.9	1.7	1.0	1.4	1.0
Jordan	0.4	0.0	0.0	0.3	0.3	0.6	0.4
Korea, Rep. of	0.0	0.1	0.2	0.1	0.8	0.6	0.4
Malaysia	0.4	0.7	1.4	1.2	1.7	2.7	1.7
Nepal	0.0	0.0	0.1	0.1	0.1	0.0	0.0
Pakistan ^c	0.0	0.3	0.8	1.2	1.1	2.3	1.2
Philippines	0.2	1.1	1.9	2.2	1.6	2.2	1.8
Sri Lanka	0.0	1.8	3.0	4.5	3.0	5.2	3.5
Thailand	0.0	0.8	0.3	0.9	0.7	0.9	0.7
<i>Caribbean and Latin America</i>							
Colombia	0.8	0.2	1.4	0.7	1.0	1.0	0.9
Costa Rica ^e	0.0	0.5	0.2	0.2	0.0	1.1	0.5
Dominican Rep.	N Ask						
Guyana	0.9	0.8	1.4	2.2	1.4	0.8	1.2
Jamaica	0.3	0.0	0.0	1.0	0.5	0.6	0.4
Mexico	N Ask						
Panama ^e	0.0	0.9	2.1	0.9	1.9	1.6	1.4
Peru	0.4	0.9	1.2	1.4	3.3	3.2	2.1

Table 4.1.9.2 Percentage of Currently Married Women Sterilized for Contraceptive Purposes, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.0	0.0	0.3	0.5	0.6	0.3
Fiji	0.4	0.5	3.4	8.0	22.7	34.3	15.8
Indonesia	0.0	0.0	0.1	0.1	0.4	0.8	0.3
Jordan	0.5	0.0	0.4	0.6	0.5	3.1	1.8
Korea, Rep. of	0.0	0.1	2.4	2.5	2.2	1.7	1.7
Malaysia	0.0	0.4	1.0	1.6	4.4	6.7	3.4
Nepal	0.0	0.0	0.1	0.1	0.0	0.2	0.1
Pakistan ^c	0.1	0.1	0.6	0.8	0.9	1.9	0.9
Philippines	0.4	0.8	3.0	6.8	7.3	5.3	4.7
Sri Lanka	0.0	0.4	2.3	9.3	14.0	17.4	9.2
Thailand	0.0	0.6	3.8	8.6	12.2	8.9	6.3
<i>Caribbean and Latin America</i>							
Colombia	0.8	0.0	1.4	5.3	5.2	6.8	4.0
Costa Rica ^e	1.6	1.4	3.7	13.8	18.2	22.2	12.3
Dominican Rep.	0.9	0.8	8.1	17.5	19.7	17.6	11.9
Guyana	0.5	0.4	1.8	5.6	7.1	19.2	8.5
Jamaica	0.7	1.7	1.9	4.7	8.7	20.3	8.1
Mexico	0.4	0.6	1.4	2.6	2.7	4.5	2.7
Panama ^e	1.7	3.0	10.5	25.6	28.8	31.8	21.2
Peru	0.0	0.2	0.9	4.4	3.2	4.3	2.8

Table 4.1.10.2 Percentage of Currently Married Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.1	0.3	0.5	0.8	0.9	0.5
Fiji	0.0	0.0	0.1	0.3	0.0	0.0	0.1
Indonesia	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Jordan	0.0	0.0	0.0	0.1	0.0	0.2	0.1
Korea, Rep. of	0.0	0.3	2.4	4.6	5.3	4.1	3.3
Malaysia	0.0	0.1	0.1	0.3	0.3	0.7	0.4
Nepal	0.1	0.3	1.0	1.8	3.0	4.5	1.6
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.2	0.1
Philippines	0.0	0.4	0.3	0.6	1.1	0.8	0.7
Sri Lanka	0.0	0.1	0.6	1.0	1.5	0.8	0.7
Thailand	0.8	0.2	2.3	4.8	4.7	1.0	2.1
<i>Caribbean and Latin America</i>							
Colombia	0.0	0.0	0.0	0.2	0.3	0.3	0.2
Costa Rica ^e	0.0	0.7	0.6	1.7	1.1	1.1	1.0
Dominican Rep.	0.0	0.0	0.0	0.0	0.0	0.2	0.1
Guyana	0.0	0.0	0.2	0.0	0.0	0.0	0.0
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.2	0.1	0.4	0.1	0.1	0.2	0.2
Panama ^e	0.0	0.0	0.6	0.4	0.3	0.4	0.4
Peru	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 4.1.11.2 Percentage of Currently Married Women Currently Using Injection, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	N Ask						
Fiji	0.0	0.3	1.0	0.3	0.2	0.1	0.3
Indonesia	0.0	0.1	0.3	0.3	0.2	0.5	0.2
Jordan	N Ask						
Korea, Rep. of	0.0	0.1	0.2	0.2	0.3	0.2	0.2
Malaysia	0.0	0.1	0.2	0.3	0.3	0.2	0.2
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Philippines	0.0	0.1	0.3	0.4	0.2	0.1	0.2
Sri Lanka	0.0	0.2	0.6	0.5	0.2	0.2	0.3
Thailand	0.3	3.3	2.4	1.3	2.1	1.6	1.9
<i>Caribbean and Latin America</i>							
Colombia	1.2	0.2	0.6	0.0	0.3	0.4	0.4
Costa Rica ^e	0.5	0.2	1.2	1.7	1.1	4.0	2.0
Dominican Rep.	0.0	0.4	0.4	0.0	0.0	0.2	0.2
Guyana	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Jamaica	0.7	2.9	9.3	5.4	10.0	8.1	6.2
Mexico	0.4	1.9	2.9	2.0	1.2	1.5	1.7
Panama ^e	0.0	1.2	1.5	0.0	0.5	0.4	0.6
Peru	0.4	0.8	1.4	1.9	1.0	0.7	1.0

Table 4.1.12.2 Percentage of Currently Married Women Currently Using Other Methods, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.0	0.0	0.2	0.3	0.5	0.2
Fiji	0.0	0.5	0.8	0.2	0.2	0.5	0.4
Indonesia	0.2	0.9	1.1	1.8	1.6	1.7	1.2
Jordan	0.0	1.8	1.1	1.6	1.6	2.8	2.0
Korea, Rep. of	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Malaysia	0.2	1.6	2.3	2.4	3.0	2.5	2.2
Nepal	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pakistan ^c	0.0	0.0	0.0	0.0	0.1	0.3	0.1
Philippines	0.0	0.0	0.1	0.1	0.0	0.1	0.1
Sri Lanka	0.0	0.0	0.3	0.0	0.0	0.0	0.1
Thailand	0.0	0.4	0.2	0.7	0.0	0.2	0.3
<i>Caribbean and Latin America</i>							
Colombia	0.0	1.0	1.9	0.5	0.6	0.6	0.8
Costa Rica ^e	0.0	0.5	0.2	0.0	0.4	0.7	0.4
Dominican Rep.	0.0	0.4	0.0	0.0	1.0	0.3	0.3
Guyana	0.7	0.4	0.2	0.0	0.0	0.5	0.4
Jamaica	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Mexico	0.4	0.0	0.2	0.1	0.4	0.0	0.2
Panama ^e	0.6	0.3	0.4	0.2	0.0	0.6	0.4
Peru	0.4	0.1	1.2	0.4	0.6	0.7	0.6

Table 4.2.1.1 Percentage of Exposed^h Women Currently Using the Pill, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.4	3.7	5.7	6.3	3.2	2.3	1.5	3.7
Fiji	9.3	14.7	13.2	9.0	9.0	6.9	3.2	10.5
Indonesia	11.1	22.9	23.9	23.9	20.3	18.8	15.4	20.8
Jordan	8.5	15.0	21.2	18.1	19.7	20.4	13.6	17.6
Korea, Rep. of	10.3	7.3	9.0	12.1	15.7	9.3	5.7	11.0
Malaysia	16.2	24.3	29.8	21.3	18.2	13.5	6.4	20.7
Nepal	0.0	0.2	0.4	1.3	0.7	0.7	0.0	0.5
Pakistan ^c	0.0	0.2	1.5	2.5	2.5	1.7	0.4	1.3
Philippines	3.6	8.1	7.6	6.6	5.5	3.5	2.5	6.0
Sri Lanka	1.4	2.8	2.5	2.4	2.2	0.6	0.4	2.0
Thailand	16.8	19.7	25.1	19.6	16.3	13.4	6.8	18.7
<i>Caribbean and Latin America</i>								
Colombia	17.3	22.0	22.0	19.2	10.7	7.6	6.8	16.2
Costa Rica	NA	46.7	36.3	29.4	16.2	12.1	4.3	27.2
Dominican Rep.	7.3	17.7	15.5	8.5	5.9	2.9	3.7	10.3
Guyana	11.5	12.2	16.8	12.9	6.8	4.2	0.9	10.6
Jamaica	14.9	17.7	22.5	13.9	9.0	6.2	3.4	13.8
Mexico	11.6	18.0	18.6	16.0	12.4	9.1	7.4	14.9
Panama	NA	33.7	32.4	19.1	12.4	8.0	2.5	20.6
Peru	4.6	5.5	9.2	7.2	4.1	2.3	1.1	5.5

Table 4.2.2.1 Percentage of Exposed^h Women Currently Using IUD, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.2	0.0	0.7	2.1	0.7	0.7	0.5
Fiji	0.6	5.4	6.6	7.1	5.9	6.9	4.8	6.0
Indonesia	1.4	5.7	7.0	10.1	11.1	11.2	10.1	7.9
Jordan	0.6	0.8	3.5	5.0	3.6	3.5	1.9	3.0
Korea, Rep. of	2.6	3.5	8.4	11.2	13.9	12.3	11.5	10.5
Malaysia	0.0	0.3	0.8	1.6	0.9	1.5	0.4	0.9
Nepal	0.0	0.1	0.0	0.1	0.1	0.0	0.6	0.1
Pakistan ^c	0.0	0.3	1.0	0.9	2.0	1.0	2.2	0.9
Philippines	0.6	2.4	3.0	3.6	3.6	3.0	3.0	3.1
Sri Lanka	7.3	7.4	5.6	7.4	5.9	6.8	1.9	6.2
Thailand	1.2	8.7	6.2	13.3	9.0	6.9	4.3	8.1
<i>Caribbean and Latin America</i>								
Colombia	5.5	8.0	13.6	14.5	12.3	7.0	3.7	10.4
Costa Rica	NA	5.8	5.9	7.5	7.6	5.2	4.8	6.3
Dominican Rep.	0.0	2.8	7.2	4.7	3.5	2.2	0.9	3.7
Guyana	1.2	7.2	9.5	10.0	5.8	2.6	3.5	6.5
Jamaica	1.0	1.0	1.7	3.7	4.2	2.1	3.4	2.3
Mexico	2.4	6.2	11.4	9.0	8.6	5.6	1.6	7.8
Panama	NA	4.2	6.1	5.8	3.5	2.8	2.0	4.5
Peru	0.6	1.8	1.7	3.5	1.2	1.5	0.4	1.8

Table 4.2.3.1 Percentage of Exposed^h Women Currently Using Other Female Scientific Methods, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.0	0.1	0.0	0.0	0.3	0.0	0.1
Fiji	0.0	0.0	0.1	0.0	0.0	0.2	0.0	0.1
Indonesia	0.0	0.0	0.2	0.2	0.1	0.0	0.0	0.1
Jordan	0.0	0.3	0.2	0.7	0.0	0.0	0.0	0.2
Korea, Rep. of	0.0	0.0	0.5	0.6	0.1	0.5	0.0	0.4
Malaysia	0.0	0.0	0.1	0.2	0.1	0.3	0.7	0.2
Nepal	N Ask							
Pakistan ^c	0.0	0.3	0.1	0.1	0.5	0.4	0.4	0.2
Philippines	0.0	0.1	0.1	0.1	0.4	0.3	0.6	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Thailand	0.0	0.2	0.0	0.2	0.2	0.0	0.0	0.1
<i>Caribbean and Latin America</i>								
Colombia	2.4	2.7	2.1	4.4	3.4	2.8	1.0	2.8
Costa Rica	NA	1.5	2.0	2.6	2.3	2.4	1.4	2.1
Dominican Rep.	1.8	2.1	2.2	2.8	2.7	0.7	0.9	2.1
Guyana	0.0	0.4	0.5	0.7	0.0	0.3	0.4	0.4
Jamaica	2.0	1.9	0.6	3.7	1.4	0.8	1.7	1.7
Mexico	0.7	1.6	2.1	3.0	1.5	0.9	2.6	1.9
Panama	NA	1.6	3.3	2.2	2.7	1.0	0.5	2.1
Peru	0.6	0.7	0.7	2.1	1.6	1.0	0.0	1.1

Table 4.2.4.1 Percentage of Exposed^h Women Currently Using Douche, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.0	0.3	0.4	0.0	0.0	0.0	0.1
Fiji	N Ask							
Indonesia	0.5	0.0	0.0	0.1	0.0	0.0	0.1	0.1
Jordan	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0
Korea, Rep. of	0.0	0.0	0.0	0.5	0.6	0.5	0.0	0.3
Malaysia	N Ask							
Nepal								
Pakistan								
Philippines	0.8	0.1	0.1	0.2	0.1	0.3	0.5	0.2
Sri Lanka	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0
Thailand	0.0	0.0	0.2	0.2	0.0	0.0	0.0	0.1
<i>Caribbean and Latin America</i>								
Colombia	1.6	0.0	0.6	1.2	0.3	0.9	0.5	0.6
Costa Rica	NA	0.7	0.4	0.0	0.3	0.3	0.5	0.4
Dominican Rep.	2.8	0.0	0.7	1.4	0.8	0.0	0.0	0.7
Guyana	3.2	1.5	1.8	3.1	2.5	3.2	0.4	2.2
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.3	0.1	0.0	0.3	0.8	0.2	0.0	0.3
Panama	NA	1.3	0.8	0.4	0.8	1.0	1.0	0.8
Peru	3.7	4.4	3.6	4.8	4.9	4.2	5.1	4.4

Table 4.2.5.1 Percentage of Exposed^h Women Currently Using Condom, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.0	1.0	0.7	0.9	0.8	1.2	0.7	0.9
Fiji	13.0	11.5	10.7	6.7	3.8	3.5	2.1	7.7
Indonesia	0.8	1.7	3.2	2.8	3.0	4.6	1.2	2.6
Jordan	0.9	1.1	1.2	2.4	3.8	2.8	2.5	2.1
Korea, Rep. of	0.0	3.3	8.1	8.0	7.8	4.8	2.5	6.7
Malaysia	0.6	2.2	4.4	4.4	4.1	2.9	1.8	3.5
Nepal	0.4	0.5	0.3	0.1	0.2	0.0	0.0	0.3
Pakistan ^c	0.0	1.4	2.3	1.7	2.0	0.9	0.8	1.4
Philippines	2.5	5.5	5.0	4.9	5.2	4.3	1.8	4.7
Sri Lanka	4.3	2.9	3.8	4.0	3.5	0.8	0.3	3.0
Thailand	0.0	0.9	0.4	0.6	0.7	0.9	0.0	0.6
<i>Caribbean and Latin America</i>								
Colombia	2.4	2.4	1.6	2.0	4.2	0.6	0.5	2.1
Costa Rica	NA	10.5	11.0	11.4	10.4	10.0	10.1	10.7
Dominican Rep.	1.8	2.1	0.7	2.8	2.0	3.6	1.9	2.0
Guyana	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0
Jamaica	12.9	12.1	8.1	5.8	5.9	2.9	2.3	7.6
Mexico	0.3	0.3	1.6	0.9	1.2	1.2	2.1	1.0
Panama	NA	2.1	1.6	1.4	1.6	1.4	0.5	1.5
Peru	0.0	1.3	0.7	2.3	1.6	1.9	0.7	1.4

Table 4.2.6.1 Percentage of Exposed^h Women Currently Using Rhythm, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.4	1.4	1.6	2.0	1.1	2.5	0.7	1.4
Fiji	1.2	2.4	2.4	4.5	2.5	3.5	2.7	3.0
Indonesia	0.0	0.5	1.7	1.6	1.7	1.0	1.4	1.2
Jordan	0.5	2.2	2.7	5.5	3.5	3.6	1.9	3.1
Korea, Rep. of	5.1	1.8	4.9	7.4	6.3	8.6	4.5	6.0
Malaysia	0.6	3.5	4.7	5.7	4.3	5.1	3.2	4.4
Nepal	N Ask							
Pakistan ^c	0.0	0.0	0.2	0.2	0.4	0.2	0.0	0.2
Philippines	5.0	7.4	11.3	12.7	13.3	12.3	9.2	11.3
Sri Lanka	3.7	5.0	9.5	13.0	12.0	12.3	10.7	10.4
Thailand	1.1	1.6	0.8	0.9	0.9	1.8	2.4	1.2
<i>Caribbean and Latin America</i>								
Colombia	3.9	7.7	4.9	5.4	6.0	7.6	7.3	6.2
Costa Rica	NA	4.1	4.9	6.8	7.4	8.6	6.3	6.2
Dominican Rep.	0.0	1.8	2.2	1.9	1.2	2.2	0.9	1.6
Guyana	2.0	4.6	3.0	3.1	5.8	1.9	0.9	3.4
Jamaica	0.0	0.2	0.3	0.7	0.3	0.8	0.0	0.4
Mexico	0.7	3.0	5.4	3.7	6.1	5.2	2.6	4.2
Panama	NA	4.7	2.5	2.6	2.7	2.8	3.4	3.1
Peru	6.3	16.2	17.6	16.1	11.9	13.5	9.3	14.3

Table 4.2.7.1 Percentage of Exposed^h Women Currently Using Withdrawal, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.7	0.8	0.5	0.3	1.0	0.3	1.1	0.6
Fiji	2.5	5.1	3.4	3.2	4.2	2.2	2.1	3.5
Indonesia	0.6	0.5	0.4	0.3	0.4	0.8	0.0	0.5
Jordan	2.4	2.7	5.5	5.7	4.9	6.4	7.0	4.9
Korea, Rep. of	0.0	2.3	2.5	3.4	4.8	4.5	0.0	3.4
Malaysia	2.3	1.5	2.2	2.2	2.4	4.2	3.2	2.5
Nepal	N Ask							
Pakistan ^c	0.0	0.1	0.0	0.5	0.1	0.0	0.0	0.1
Philippines	9.2	11.7	12.3	12.2	13.0	13.3	13.2	12.5
Sri Lanka	0.6	2.3	2.5	2.5	1.3	1.4	1.0	1.9
Thailand	1.2	0.9	1.5	0.8	0.7	1.8	2.8	1.2
<i>Caribbean and Latin America</i>								
Colombia	2.4	4.6	4.7	7.9	3.9	9.5	6.3	5.8
Costa Rica	NA	5.6	6.1	4.7	6.1	6.2	4.3	5.6
Dominican Rep.	4.6	5.0	5.1	3.8	5.5	6.5	2.8	4.9
Guyana	1.2	1.8	0.9	0.9	1.0	0.6	1.8	1.2
Jamaica	1.0	1.7	2.0	2.4	1.4	1.2	1.7	1.7
Mexico	2.7	3.8	4.2	6.0	5.8	6.6	6.3	5.0
Panama	NA	2.6	3.5	3.4	4.3	3.5	4.9	3.6
Peru	3.4	3.5	4.8	4.9	4.7	4.2	3.2	4.3

Table 4.2.8.1 Percentage of Exposed^h Women Currently Using Abstinence, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	1.2	1.5	0.5	1.9	3.0	2.0	1.6	1.5
Fiji	0.0	0.3	0.2	0.3	0.2	1.2	0.0	0.3
Indonesia	0.5	1.0	0.8	1.3	1.8	2.9	4.0	1.4
Jordan	0.0	0.5	0.6	0.5	0.9	0.0	2.0	0.6
Korea, Rep. of	0.0	0.3	0.0	0.1	1.4	0.7	0.6	0.5
Malaysia	0.0	0.9	0.8	1.6	2.3	4.9	7.1	2.2
Nepal	0.0	0.1	0.0	0.0	0.3	0.0	0.0	0.1
Pakistan ^c	0.1	0.3	1.1	1.8	2.1	4.2	5.0	1.6
Philippines	0.5	1.7	2.6	2.5	2.6	2.9	2.5	2.4
Sri Lanka	0.4	2.1	2.0	3.5	4.3	7.9	14.2	4.5
Thailand	0.6	0.7	0.4	0.8	1.3	1.2	2.9	0.9
<i>Caribbean and Latin America</i>								
Colombia	0.0	1.2	1.2	1.5	0.8	1.3	1.0	1.1
Costa Rica	NA	0.2	0.6	0.0	0.5	1.0	2.4	0.6
Dominican Rep.	N Ask							
Guyana	2.0	1.7	1.1	1.3	1.0	0.6	2.2	1.4
Jamaica	0.0	0.5	0.0	0.3	1.4	0.0	1.1	0.5
Mexico	N Ask							
Panama	NA	2.3	1.6	2.0	1.3	1.4	1.5	1.7
Peru	1.1	1.4	2.7	1.9	2.4	5.6	4.4	2.8

Table 4.2.9.1 Percentage of Exposed^h Women Sterilized for Contraceptive Purposes, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.1	0.1	0.3	0.7	1.1	0.8	0.3	0.4
Fiji	0.0	1.2	12.0	25.5	32.1	36.8	43.9	20.1
Indonesia	0.0	0.0	0.1	0.4	1.0	0.7	1.0	0.4
Jordan	0.0	0.0	0.6	2.3	5.0	6.6	7.6	2.6
Korea, Rep. of	0.0	0.0	0.4	3.7	2.8	2.7	6.4	2.3
Malaysia	0.0	0.4	1.5	5.7	7.0	7.3	7.5	4.4
Nepal	0.0	0.1	0.0	0.0	0.5	0.0	0.0	0.1
Pakistan ^c	0.0	0.0	0.5	0.9	2.5	2.6	7.8	1.3
Philippines	0.0	1.9	5.5	7.6	8.4	7.0	6.2	6.2
Sri Lanka	0.6	2.2	9.3	15.2	17.2	16.0	13.0	12.0
Thailand	0.6	1.9	7.7	10.7	12.0	13.5	15.0	8.6
<i>Caribbean and Latin America</i>								
Colombia	0.0	1.0	2.7	7.6	9.2	7.0	4.2	4.9
Costa Rica	NA	0.7	8.0	15.6	26.1	24.5	22.6	14.9
Dominican Rep.	0.0	5.3	17.0	25.0	25.5	15.8	13.1	15.6
Guyana	0.0	1.1	4.7	14.7	20.3	25.4	23.8	11.4
Jamaica	0.0	0.7	3.7	16.0	19.8	23.7	27.1	11.4
Mexico	0.3	0.6	1.6	4.4	6.1	8.2	9.5	3.7
Panama	NA	1.0	16.2	27.9	36.1	41.5	47.5	25.6
Peru	0.0	0.2	1.6	5.1	7.5	3.8	5.6	3.6

Table 4.2.10.1 Percentage of Exposed^h Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.3	0.4	0.5	3.0	0.9	0.5	0.6
Fiji	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.1
Indonesia	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0
Jordan	0.0	0.0	0.0	0.0	0.1	0.2	2.0	0.2
Korea, Rep. of	0.0	0.3	1.7	3.1	6.7	7.5	13.4	4.3
Malaysia	0.0	0.0	0.2	0.2	0.8	1.0	1.4	0.5
Nepal	0.0	0.3	1.9	3.9	4.0	3.4	5.0	2.0
Pakistan ^c	0.0	0.0	0.0	0.1	0.3	0.3	0.0	0.1
Philippines	0.0	0.4	0.8	1.1	1.3	0.7	0.9	0.9
Sri Lanka	0.0	0.2	0.9	1.5	1.2	1.3	0.0	0.9
Thailand	0.0	1.1	2.3	2.0	6.0	4.0	4.3	2.9
<i>Caribbean and Latin America</i>								
Colombia	0.0	0.2	0.4	0.2	0.0	0.3	0.0	0.2
Costa Rica	NA	0.0	1.6	2.3	0.8	0.7	1.4	1.2
Dominican Rep.	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.1
Guyana	0.0	0.0	0.0	0.2	0.3	0.0	0.0	0.1
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.0	0.0	0.1	0.4	0.8	0.2	0.0	0.3
Panama	NA	0.3	0.0	1.0	0.5	0.3	0.5	0.4
Peru	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 4.2.11.1 Percentage of Exposed^h Women Currently Using Injection, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	N Ask							
Fiji	1.2	0.9	0.1	0.4	0.0	0.2	0.0	0.4
Indonesia	0.2	0.3	0.6	0.4	0.3	0.1	0.6	0.3
Jordan	N Ask							
Korea, Rep. of	0.0	0.0	0.3	0.3	0.3	0.2	0.0	0.3
Malaysia	0.0	0.4	0.3	0.4	0.2	0.2	0.0	0.3
Nepal	N Ask							
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Philippines	0.4	0.2	0.4	0.2	0.2	0.0	0.0	0.2
Sri Lanka	0.0	0.9	0.1	0.7	0.3	0.1	0.5	0.4
Thailand	1.2	2.7	3.3	1.9	3.1	2.5	1.7	2.6
<i>Caribbean and Latin America</i>								
Colombia	1.6	0.5	0.2	0.5	1.0	0.3	0.0	0.5
Costa Rica	NA	0.7	1.8	1.4	2.3	4.8	5.8	2.4
Dominican Rep.	0.9	0.4	0.0	0.0	0.0	0.0	0.9	0.2
Guyana	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Jamaica	5.5	9.7	10.7	10.5	5.9	1.2	0.6	7.2
Mexico	1.4	2.1	2.6	2.9	2.6	1.9	0.0	2.3
Panama	NA	1.8	1.0	0.2	0.5	0.3	0.5	0.8
Peru	2.4	2.4	1.9	1.1	1.0	0.2	0.7	1.4

Table 4.2.12.1 Percentage of Exposed^h Women Currently Using Other Methods, by Current Age

Country	Current Age							Total
	15-19	20-24	25-29	30-34	35-39	40-44	45-49	
<i>Asia and Pacific</i>								
Bangladesh	0.0	0.0	0.4	0.5	0.4	0.6	0.0	0.2
Fiji	1.2	0.3	0.6	0.4	0.8	0.5	0.0	0.5
Indonesia	1.1	0.8	2.1	1.9	2.2	2.3	1.9	1.7
Jordan	0.5	2.6	1.8	4.0	4.5	4.1	2.9	3.0
Korea, Rep. of	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0
Malaysia	1.7	3.6	3.4	2.2	2.9	2.2	2.9	2.8
Nepal	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pakistan ^c	0.0	0.0	0.3	0.0	0.4	0.0	0.8	0.2
Philippines	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.1
Sri Lanka	0.0	0.0	0.0	0.2	0.0	0.2	0.0	0.1
Thailand	0.0	0.7	0.3	0.8	0.0	0.0	0.0	0.3
<i>Caribbean and Latin America</i>								
Colombia	0.8	1.5	1.6	1.0	1.0	0.0	0.0	1.0
Costa Rica	NA	0.5	0.6	0.2	0.3	1.0	0.0	0.5
Dominican Rep.	0.9	0.0	0.4	0.0	0.0	1.4	0.9	0.4
Guyana	1.6	0.0	0.4	0.7	0.3	0.0	0.9	0.4
Jamaica	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.1
Mexico	0.7	0.1	0.3	0.0	0.3	0.0	0.5	0.2
Panama	NA	0.3	0.8	0.4	0.5	0.3	0.0	0.4
Peru	1.0	1.0	0.9	0.8	0.7	0.8	0.4	0.8

Table 4.2.1.2 Percentage of Exposed^h Women Currently Using the Pill, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.4	2.1	3.8	4.0	4.1	5.5	3.7
Fiji	2.8	14.1	15.5	15.4	7.8	7.7	10.5
Indonesia	3.2	19.3	25.8	26.0	21.3	23.5	20.8
Jordan	2.4	13.6	20.6	18.2	18.3	19.3	17.6
Korea, Rep. of	7.0	5.0	10.2	12.3	13.6	12.2	11.0
Malaysia	2.4	22.8	23.3	23.3	23.3	19.7	20.7
Nepal	0.0	0.4	0.0	0.6	0.6	1.8	0.5
Pakistan ^c	0.0	0.4	0.3	1.2	1.9	2.7	1.3
Philippines	0.3	6.8	8.2	7.0	6.0	4.9	6.0
Sri Lanka	0.5	2.1	2.7	2.4	2.4	1.6	2.0
Thailand	6.4	24.0	23.7	18.8	16.0	16.2	18.7
<i>Caribbean and Latin America</i>							
Colombia	9.4	21.6	18.9	19.4	16.0	12.6	16.2
Costa Rica ^e	16.8	42.5	37.0	24.9	24.3	16.9	27.2
Dominican Rep.	6.1	16.8	16.2	12.0	7.8	6.6	10.3
Guyana	9.0	15.1	16.3	11.9	10.9	6.4	10.6
Jamaica	10.9	17.2	20.3	18.6	11.2	7.8	13.8
Mexico	5.5	16.8	20.6	16.6	17.5	11.7	14.9
Panama ^e	15.4	34.6	32.7	23.0	18.6	10.1	20.6
Peru	3.9	5.9	8.4	6.7	5.0	3.5	5.5

Table 4.2.2.2 Percentage of Exposed^h Women Currently Using IUD, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.2	0.0	0.5	0.3	1.1	0.5
Fiji	0.0	2.2	6.2	5.7	11.4	7.2	6.0
Indonesia	0.0	3.1	7.2	10.0	11.3	13.5	7.9
Jordan	0.0	1.4	3.7	1.4	3.2	3.8	3.0
Korea, Rep. of	0.0	2.0	10.5	11.5	14.0	13.2	10.5
Malaysia	0.4	0.7	1.1	1.3	0.7	0.9	0.9
Nepal	0.0	0.0	0.1	0.1	0.0	0.2	0.1
Pakistan ^c	0.1	0.3	0.4	0.8	0.8	1.9	0.9
Philippines	0.1	1.6	3.3	3.6	4.5	3.0	3.1
Sri Lanka	0.0	4.1	7.4	7.5	8.2	6.5	6.2
Thailand	0.0	4.0	8.7	12.1	10.4	8.8	8.1
<i>Caribbean and Latin America</i>							
Colombia	0.0	8.4	13.0	14.1	15.3	8.4	10.4
Costa Rica ^e	2.1	4.8	5.9	7.7	7.0	6.9	6.3
Dominican Rep.	0.0	1.1	4.9	5.8	4.2	4.1	3.7
Guyana	0.0	2.1	7.6	8.3	8.0	9.1	6.5
Jamaica	1.2	0.3	1.8	1.9	4.3	3.9	2.3
Mexico	0.4	6.7	9.2	9.8	9.3	7.3	7.8
Panama ^e	0.0	4.5	6.3	4.5	5.7	3.7	4.5
Peru	0.0	2.0	3.4	1.2	1.8	1.2	1.8

Table 4.2.3.2 Percentage of Exposed^h Women Currently Using Other Female Scientific Methods, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.0	0.0	0.0	0.1	0.1	0.1
Fiji	0.0	0.2	0.0	0.2	0.0	0.0	0.1
Indonesia	0.0	0.1	0.0	0.0	0.1	0.2	0.1
Jordan	0.0	0.7	0.4	0.4	0.0	0.2	0.2
Korea, Rep. of	0.0	0.6	0.1	0.6	0.4	0.3	0.4
Malaysia	0.0	0.0	0.1	0.4	0.3	0.1	0.2
Nepal	N Ask						
Pakistan ^c	0.0	0.3	0.0	0.4	0.0	0.4	0.2
Philippines	0.0	0.0	0.1	0.3	0.2	0.3	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Thailand	0.0	0.0	0.2	0.0	0.5	0.0	0.1
<i>Caribbean and Latin America</i>							
Colombia	1.4	2.5	4.9	2.3	3.4	2.3	2.8
Costa Rica ^e	2.1	1.1	3.3	4.0	1.2	1.1	2.1
Dominican Rep.	2.3	1.6	2.9	1.4	0.6	2.7	2.1
Guyana	0.0	0.0	0.3	0.3	1.0	0.5	0.4
Jamaica	2.8	1.4	1.8	2.3	0.0	1.7	1.7
Mexico	2.1	1.0	3.0	2.8	2.6	1.0	1.9
Panama ^e	1.9	3.3	2.8	4.0	0.6	1.1	2.1
Peru	1.7	0.4	1.4	1.3	1.1	1.0	1.1

Table 4.2.4.2 Percentage of Exposed^h Women Currently Using Douche, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.0	0.2	0.0	0.4	0.1	0.1
Fiji	N Ask						
Indonesia	0.0	0.3	0.1	0.0	0.0	0.1	0.1
Jordan	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Korea, Rep. of	0.0	0.0	0.4	0.5	0.4	0.3	0.3
Malaysia	N Ask						
Nepal							
Pakistan							
Philippines	0.0	0.5	0.2	0.0	0.4	0.2	0.2
Sri Lanka	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Thailand	0.0	0.0	0.2	0.0	0.3	0.0	0.1
<i>Caribbean and Latin America</i>							
Colombia	0.7	0.0	1.2	1.4	0.0	0.5	0.6
Costa Rica ^e	0.0	0.3	0.4	0.6	0.4	0.3	0.4
Dominican Rep.	1.5	0.0	1.0	0.5	1.8	0.4	0.7
Guyana	2.1	0.8	1.6	3.0	3.5	2.4	2.2
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.0	0.2	0.5	0.3	0.2	0.3	0.3
Panama ^e	1.9	0.7	0.8	0.8	1.3	0.6	0.8
Peru	3.5	3.4	4.4	3.9	4.4	5.1	4.4

Table 4.2.5.2 Percentage of Exposed^h Women Currently Using the Condom, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	1.0	1.1	0.5	1.0	1.0	1.0	0.9
Fiji	3.4	10.1	10.9	11.5	7.6	4.8	7.7
Indonesia	0.0	1.0	1.9	4.0	3.4	4.4	2.6
Jordan	0.0	2.7	2.2	3.0	1.8	2.1	2.1
Korea, Rep. of	2.3	4.0	9.3	8.2	7.3	5.2	6.7
Malaysia	2.4	4.2	4.3	4.5	3.0	2.8	3.5
Nepal	0.1	0.1	0.8	0.0	0.2	0.5	0.3
Pakistan ^c	0.0	0.8	1.3	1.7	1.8	2.2	1.4
Philippines	0.1	4.8	5.4	4.4	5.1	4.8	4.7
Sri Lanka	1.6	3.9	4.6	4.2	3.0	1.4	3.0
Thailand	0.5	0.4	1.1	0.2	1.2	0.3	0.6
<i>Caribbean and Latin America</i>							
Colombia	2.9	2.8	2.2	2.5	1.5	1.6	2.1
Costa Rica ^e	7.4	10.8	15.3	12.9	10.7	7.1	10.7
Dominican Rep.	0.8	2.7	2.5	1.4	3.0	1.8	2.0
Guyana	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Jamaica	11.3	8.3	11.2	6.2	8.5	3.9	7.6
Mexico	0.0	0.4	0.9	1.8	1.1	1.2	1.0
Panama ^e	1.9	2.6	1.8	1.3	2.2	0.8	1.5
Peru	0.0	0.7	1.5	2.5	1.3	1.2	1.4

Table 4.2.6.2 Percentage of Exposed^h Women Currently Using Rhythm, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.4	1.1	0.6	2.5	1.6	1.6	1.4
Fiji	1.5	1.2	4.0	3.7	4.2	2.8	3.0
Indonesia	0.3	0.6	1.4	0.9	1.8	1.8	1.2
Jordan	0.6	2.1	6.6	3.0	6.3	2.2	3.1
Korea, Rep. of	6.4	2.4	6.6	7.7	7.2	4.8	6.0
Malaysia	2.0	3.7	6.3	6.7	5.2	3.0	4.4
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.0	0.2	1.0	0.1	0.2
Philippines	2.3	9.1	12.1	13.7	12.7	11.0	11.3
Sri Lanka	1.7	9.9	15.6	12.3	15.1	7.1	10.4
Thailand	0.6	1.8	1.5	0.7	1.2	0.9	1.2
<i>Caribbean and Latin America</i>							
Colombia	3.6	9.4	8.1	3.9	9.0	4.6	6.2
Costa Rica ^e	4.2	5.1	7.8	7.4	8.2	4.6	6.2
Dominican Rep.	2.3	1.1	1.0	3.4	0.6	1.4	1.6
Guyana	4.5	2.1	4.2	3.6	3.8	2.9	3.4
Jamaica	0.4	0.0	0.0	1.6	1.1	0.0	0.4
Mexico	1.7	3.6	5.6	5.5	4.8	3.5	4.2
Panama ^e	2.9	3.7	4.8	3.4	2.2	2.1	3.1
Peru	3.9	18.4	19.0	17.2	13.6	10.0	14.3

Table 4.2.7.2 Percentage of Exposed^h Women Currently Using Withdrawal, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.0	0.9	0.9	0.7	0.8	0.5	0.6
Fiji	1.2	5.7	5.3	4.0	3.6	2.3	3.5
Indonesia	0.0	0.5	0.6	0.3	0.6	0.6	0.5
Jordan	2.4	3.4	1.9	6.4	3.6	6.1	4.9
Korea, Rep. of	0.0	1.8	3.3	4.4	3.3	4.0	3.4
Malaysia	2.0	1.5	1.8	2.4	3.5	2.8	2.5
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.4	0.0	0.2	0.1	0.1
Philippines	1.5	9.5	12.8	13.4	12.6	13.9	12.5
Sri Lanka	0.7	2.5	2.4	2.8	1.2	1.5	1.9
Thailand	0.6	1.6	0.6	0.9	1.8	1.5	1.2
<i>Caribbean and Latin America</i>							
Colombia	0.7	3.4	5.2	5.4	5.2	8.2	5.8
Costa Rica ^e	2.1	6.5	5.9	5.4	4.1	5.9	5.6
Dominican Rep.	0.8	4.3	4.9	6.3	6.6	4.9	4.9
Guyana	1.5	1.5	1.3	1.4	1.0	0.8	1.2
Jamaica	2.0	1.1	1.5	0.4	3.2	2.0	1.7
Mexico	0.0	5.5	2.9	3.8	5.0	6.9	5.0
Panama ^e	2.9	1.9	2.8	3.2	3.2	5.0	3.6
Peru	0.0	3.0	4.9	4.9	5.7	4.0	4.3

Table 4.2.8.2 Percentage of Exposed^h Women Currently Using Abstinence, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	1.2	1.5	1.1	1.3	2.4	1.5
Fiji	0.0	0.2	0.4	0.2	0.0	0.6	0.3
Indonesia	0.2	1.1	1.2	2.2	1.3	2.0	1.4
Jordan	0.7	0.0	0.0	0.4	0.4	0.9	0.6
Korea, Rep. of	0.0	0.2	0.3	0.1	0.9	0.9	0.5
Malaysia	0.8	0.9	1.7	1.4	2.0	3.4	2.2
Nepal	0.0	0.0	0.1	0.1	0.1	0.0	0.1
Pakistan ^c	0.0	0.4	1.1	1.7	1.6	3.2	1.6
Philippines	0.4	1.6	2.3	2.7	1.9	3.0	2.4
Sri Lanka	0.0	2.2	3.8	5.4	3.7	7.2	4.5
Thailand	0.0	0.9	0.4	1.2	1.0	1.3	0.9
<i>Caribbean and Latin America</i>							
Colombia	1.4	0.3	1.7	0.8	1.1	1.2	1.1
Costa Rica ^e	0.0	0.6	0.2	0.3	0.0	1.4	0.6
Dominican Rep.	N Ask						
Guyana	1.2	1.0	1.6	2.5	1.6	0.9	1.4
Jamaica	0.4	0.0	0.0	1.2	0.5	0.7	0.5
Mexico	N Ask						
Panama ^e	0.0	1.1	2.5	1.1	2.2	1.9	1.7
Peru	0.9	1.2	1.4	1.7	4.1	4.2	2.8

Table 4.2.9.2 Percentage of Exposed^h Women Sterilized for Contraceptive Purposes, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.0	0.0	0.4	0.6	0.8	0.4
Fiji	0.6	0.6	4.4	9.5	26.6	42.3	20.1
Indonesia	0.0	0.0	0.1	0.1	0.5	1.2	0.4
Jordan	0.9	0.0	0.5	0.8	0.7	4.6	2.6
Korea, Rep.of	0.0	0.2	2.9	3.0	2.7	2.4	2.3
Malaysia	0.0	0.6	1.3	2.0	5.2	8.4	4.4
Nepal	0.0	0.0	0.1	0.2	0.0	0.3	0.1
Pakistan ^c	0.1	0.1	0.8	1.2	1.4	2.6	1.3
Philippines	0.7	1.0	3.8	8.3	9.0	7.3	6.2
Sri Lanka	0.0	0.4	2.8	11.3	17.4	24.1	12.0
Thailand	0.0	0.7	4.8	10.8	16.6	13.5	8.6
<i>Caribbean and Latin America</i>							
Colombia	1.4	0.0	1.7	6.2	6.0	7.9	4.9
Costa Rica ^e	3.2	1.7	4.1	16.0	21.4	26.8	14.9
Dominican Rep.	1.5	1.1	10.3	20.7	24.1	22.1	15.6
Guyana	2.7	2.6	3.4	7.8	8.7	23.4	11.4
Jamaica	3.6	4.2	3.9	7.8	13.3	24.4	11.4
Mexico	0.8	0.8	1.8	3.3	3.4	6.2	3.7
Panama ^e	2.9	3.7	12.4	31.4	33.8	36.4	25.6
Peru	0.0	0.2	1.1	5.5	4.0	5.6	3.6

Table 4.2.10.2 Percentage of Exposed^h Women Whose Husbands Have Been Sterilized for Contraceptive Purposes, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.2	0.2	0.3	0.5	1.0	1.0	0.6
Fiji	0.0	0.0	0.0	0.4	0.0	0.0	0.1
Indonesia	0.0	0.0	0.0	0.1	0.0	0.1	0.0
Jordan	0.0	0.0	0.0	0.2	0.0	0.3	0.2
Korea, Rep. of	0.0	0.4	2.9	5.3	6.5	5.7	4.3
Malaysia	0.0	0.2	0.1	0.4	0.3	0.9	0.5
Nepal	0.1	0.4	1.2	2.2	3.9	6.5	2.0
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.3	0.1
Philippines	0.0	0.5	0.4	0.7	1.3	1.1	0.9
Sri Lanka	0.0	0.2	0.8	1.2	1.9	1.1	0.9
Thailand	1.3	0.2	2.9	6.0	6.4	1.4	2.9
<i>Caribbean and Latin America</i>							
Colombia	0.0	0.0	0.0	0.3	0.4	0.4	0.2
Costa Rica ^e	0.0	0.8	0.7	2.0	1.2	1.4	1.2
Dominican Rep.	0.0	0.0	0.0	0.0	0.0	0.2	0.1
Guyana	0.0	0.0	0.3	0.0	0.0	0.1	0.1
Jamaica	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mexico	0.4	0.2	0.5	0.2	0.2	0.3	0.3
Panama ^e	0.0	0.0	0.8	0.5	0.3	0.5	0.4
Peru	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 4.2.11.2 Percentage of Exposed^h Women Currently Using Injection, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	N Ask						
Fiji	0.0	0.4	1.3	0.4	0.2	0.1	0.4
Indonesia	0.0	0.1	0.4	0.4	0.3	0.7	0.3
Jordan	N Ask						
Korea, Rep. of	0.0	0.2	0.3	0.2	0.4	0.2	0.3
Malaysia	0.0	0.2	0.3	0.4	0.3	0.3	0.3
Nepal	N Ask						
Pakistan ^c	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Philippines	0.0	0.1	0.3	0.5	0.3	0.1	0.2
Sri Lanka	0.0	0.3	0.8	0.6	0.2	0.3	0.4
Thailand	0.6	4.1	3.0	1.6	2.9	2.4	2.6
<i>Caribbean and Latin America</i>							
Colombia	2.2	0.3	0.7	0.0	0.4	0.5	0.5
Costa Rica ^e	1.1	0.3	1.3	2.0	1.2	4.8	2.4
Dominican Rep.	0.0	0.5	0.5	0.0	0.0	0.2	0.2
Guyana	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Jamaica	0.8	3.3	10.6	6.2	11.7	9.2	7.2
Mexico	0.8	2.6	3.6	2.4	1.5	2.0	2.3
Panama ^e	0.0	1.5	1.8	0.0	0.6	0.5	0.8
Peru	0.9	1.0	1.8	2.4	1.3	0.8	1.4

Table 4.2.12.2 Percentage of Exposed^h Women Currently Using Other Methods, by Number of Living Children

Country	Number of Living Children						Total
	None	1	2	3	4	5+	
<i>Asia and Pacific</i>							
Bangladesh	0.1	0.0	0.0	0.2	0.3	0.6	0.2
Fiji	0.0	0.6	1.1	0.2	0.2	0.6	0.5
Indonesia	0.3	1.2	1.5	2.3	2.2	2.5	1.7
Jordan	0.0	2.8	1.6	2.3	2.3	4.1	3.0
Korea, Rep. of	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Malaysia	0.4	2.2	2.8	2.9	3.5	3.1	2.8
Nepal	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pakistan ^c	0.0	0.0	0.0	0.0	0.2	0.4	0.2
Philippines	0.0	0.0	0.1	0.1	0.0	0.1	0.1
Sri Lanka	0.0	0.0	0.4	0.0	0.0	0.0	0.1
Thailand	0.0	0.5	0.2	0.9	0.0	0.3	0.3
<i>Caribbean and Latin America</i>							
Colombia	0.0	1.3	2.2	0.6	0.7	0.7	1.0
Costa Rica ^e	0.0	0.6	0.2	0.0	0.4	0.8	0.5
Dominican Rep.	0.0	0.5	0.0	0.0	1.2	0.4	0.4
Guyana	0.9	0.5	0.3	0.0	0.0	0.6	0.4
Jamaica	0.0	0.0	0.0	0.0	0.0	0.2	0.1
Mexico	0.8	0.0	0.3	0.2	0.6	0.1	0.2
Panama ^e	1.0	0.4	0.5	0.3	0.0	0.6	0.4
Peru	0.9	0.2	1.5	0.5	0.8	0.9	0.8

